

FORM PTO-1390
(REV 11-98)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

015280-310100US

U.S. APPLICATION NO. (if known, see 37 CFR 1.51)

09/462682

INTERNATIONAL APPLICATION NO.
PCT/US98/14341INTERNATIONAL FILING DATE
July 10, 1998PRIORITY DATE CLAIMED
July 11, 1997

TITLE OF INVENTION

PSEUDOMONAS EXOTOXIN A-LIKE CHIMERIC IMMUNOGENS

APPLICANT(S) FOR DO/EO/US

DAVID J. FITZGERALD

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau)
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:
Copy of Written Opinion of 5/25/99, Response

ATTORNEY'S DOCKET NUMBER
015280-310100US

Form PTO-1390 (REV 11-98) page 2 of 2

PATENT

Attorney Docket No. 015280-310100PC
DHHS Ref. No.: E-057-97/0

PSEUDOMONAS EXOTOXIN A-LIKE CHIMERIC IMMUNOGENS**CROSS-REFERENCE TO RELATED APPLICATION**

This application claims the benefit of the filing date of co-pending application 60/052,375, filed July 11, 1997, the content of which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

This invention is directed to the fields of chimeric proteins and immunology.

Immunization against infectious disease has been one of the great achievements of modern medicine. Vaccines can be useful only if the vaccine, itself, is not significantly pathogenic. Many vaccines are produced by inactivating the pathogen. For example, hepatitis vaccines can be made by heating the virus and treating it with formaldehyde. Other vaccines, for example certain polio vaccines, are produced by attenuating a live pathogen. However, there is concern about producing attenuated vaccines for certain infectious agents whose pathology is not fully understood, such as HIV.

Molecular biology has enabled the production of subunit vaccines; vaccines in which the immunogen is a fragment or subunit of a parent protein or complex.

Envelope proteins of HIV-1, such as gp120, are being evaluated as subunit vaccines. Several studies have suggested that antibodies to the V3 loop region of gp120 provide protection through virus neutralization. (Emeni, E.A., *et al.*, 1992, *Nature* 355, 728-30; Javaherian, K., *et al.*, 1989, *Proc Natl Acad Sci USA* 86, 6768-72; Steimer, K.S., *et al.*, 1991, *Science* 254, 105-8; Wang, C.Y., *et al.*, 1991, *Science* 254, 285-8.)

However, subunit vaccines may not be complex enough to generate an appropriate immune response. Also, when the pathogen is highly mutable, as is HIV, subunit vaccines that elicit strain-specific immunity may not be effective in providing global protection. Furthermore, the injection of inactive virus or even the envelope

protein itself has the potential to produce a mixture of neutralizing and so-called "enhancing" antibodies. (Toth, F.D., *et al.*, 1994, *Clin Exp Immunol* 96, 389-94; Eaton, A.M., *et al.*, 1994, *Aids Res Hum Retroviruses* 10, 13-8; Mitchell, W.M., *et al.*, 1995, *Aids* 9, 27-34; Montefiori, D.C., *et al.*, 1996, *J Infect Dis* 173, 60-7.)

5 The immunogenicity of subunit vaccines is sometimes increased by coupling the subunit to a carrier protein to create a conjugate vaccine. One such carrier protein is *Pseudomonas* exotoxin A ("PE"). Investigators covalently linked a non-immunogenic O-polysaccharide derived from lipopolysaccharide ("LPS") to PE. The resulting conjugate vaccine elicited an immune response against both LPS and PE. (S.J. Cryz, Jr. *et al.* (1987) *J. Clin. Invest.*, 80:51-56 and S.J. Cryz, Jr. *et al.* (1990) *J. Infectious Diseases*, 163:1040-1045.) In another study, investigators were able to evoke an immune response against a *Plasmodium falciparum* antigen by coupling it through a spacer to PE. (J.U. Que *et al.* (1988) *Infection and Immunity*, 56:2645-49.) In a third study, investigators detoxified PE and chemically cross-linked it with principle neutralizing domain ("PND") peptides of HIV-1. The conjugate vaccine elicited the production of antibodies that recognized PND peptide and neutralized the homologous strain, HIV-1_{MN}. (S.J. Cryz, Jr. *et al.* (1995) *Vaccine*, 13:66-71.)

15
20 Chimeric proteins containing components of HIV-1 have been constructed and their immunogenic properties evaluated. These include: a poliovirus antigen containing an epitope of the gp41 transmembrane glycoprotein from HIV-1 (Evans, D.J., *et al.*, 1989, *Nature* 339, 385-8), a mucin protein containing multiple copies of the V3 loop (Fontenot, J.D., *et al.*, 1995, *Proc Natl Acad Sci USA*, 92, 315-9) a genetically modified cholera B chain with V3 loop sequences (Backstrom, M., *et al.*, 1994, *Gene* 149, 211-7) and a chemically detoxified PE-V3 loop peptide conjugate (Cryz, S., Jr., *et al.*, 1995, *Vaccine* 13, 67-71).

25
30 The third variable (V3) loop of the envelope protein, gp120, contains the principal neutralizing domain of HIV-1. (Emini, E.A., *et al.*, 1992, *Nature* 355, 728-30; Javaherian, K., *et al.*, 1989, *Proc Natl Acad Sci USA* 86, 6768-72; Rusche, J.R., *et al.*, [published errata appear in *Proc Natl Acad Sci USA* 22, 8697 1988, and *Proc Natl Acad Sci USA* 5, 1667 1989, J; *Proc Natl Acad Sci USA* 85, 3198-202 1988.) Although V3 loops vary considerably amongst the various HIV-1 strains (Berman, P.W., *et al.*, 1990, *Nature* 345, 622-5) specific antibodies to this region have been shown to neutralize infectivity of the virus and to prevent viral cell fusion *in vitro* (Kovacs, J.A.,

et al. 1993, *J. Clin Invest* 92, 919-28). Further, systemic immunization with a recombinant form of gp120 appears sufficient to protect chimpanzees from infection by HIV-1 systemic challenge. White-Scharf, M.E., *et al.*, 1993, *Virology* 192, 197-206.

HIV frequently gains entry to the body at mucosal surfaces. However, presently available HIV immunogens are not known to elicit a secretory immune response, which would inhibit viral access through the mucosa.

The development of a stable vaccine that could elicit both humoral and cellular responses, including mucosal immunity, and be flexible enough to incorporate sequences from many strains of an infectious agent, such as HIV-1, would be desirable.

SUMMARY OF THE INVENTION

Pseudomonas exotoxin A-like ("PE-like") chimeric immunogens in which a non-native epitope is inserted into the Ib domain are useful to elicit humoral, cell-mediated and secretory immune responses against the non-native epitope. In particular, the non-native epitope can be the V3 loop of the gp120 protein of HIV. Such chimeras are useful in vaccines against HIV infection.

PE chimeric immunogens offer several advantages. First, they can be made by wholly recombinant means. This eliminates the need to attach the epitope to PE by chemical cross-linking and to chemically inactivate the exotoxin. Recombinant technology also allows one to make a chimeric "cassette" having an insertion site for the non-native epitope of choice at the Ib domain location. This allows one to quickly insert existing variants of an epitope, or new variants of rapidly evolving epitopes. This enables production of vaccines that include a cocktail of different immunogens.

Second, *Pseudomonas* exotoxin can be engineered to alter the function of its domains, thereby providing a variety of activities. For example, by replacing the native cell binding domain of *Pseudomonas* exotoxin A (domain Ia) with a ligand for a particular cell receptor, one can target the chimera to bind to the particular cell type.

Third, because the Ib domain includes a cysteine-cysteine loop, epitopes that are so constrained in nature can be presented in near-native conformation. This assists in provoking an immune response against the native antigen. For example, a turn-turn-helix motif is evident with circular (constrained by a disulfide bond) but not linear peptides. (Ogata, M., *et al.*, 1990, *Biol Chem* 265, 20678-85.) Also, circular

peptides are recognized more readily by anti-V3 loop monoclonal antibodies than linear ones. (Catasti, P., *et. al.*, 1995, *J Biol Chem* 270, 2224-32.)

Fourth, the chimeras of this invention can be used to elicit a humoral, a cell-mediated or a secretory immune response. *Pseudomonas* exotoxin has been reported to act as a "superantigen," binding directly to MHC Class II molecules without prior processing in the antigen presenting cell. P.K. Legaard et al. (1991) *Cellular Immunology* 135:372-382. This promotes an MHC Class II-mediated immune response against cells bearing proteins containing the non-native epitope. Also, upon binding to a cell surface receptor, chimeric *Pseudomonas* exotoxins translocate into the cytosol. This makes possible an MHC Class I-dependent immune response against cells bearing the non-native epitope on their surface. This aspect is particularly advantageous because normally the immune system mounts an MHC Class I-dependent immune response only against proteins made by the cell. Also, by directing the chimera to a mucosal surface, one can elicit a secretory immune response involving IgA.

In one aspect, this invention provides a non-toxic *Pseudomonas* exotoxin A-like ("PE-like") chimeric immunogen comprising: (1) a cell recognition domain of between 10 and 1500 amino acids that binds to a cell surface receptor; (2) a translocation domain comprising an amino acid sequence substantially identical to a sequence of PE domain II sufficient to effect translocation to a cell cytosol; (3) a non-native epitope domain comprising an amino acid sequence of between 5 and 1500 amino acids that comprises a non-native epitope; and, optionally, (4) an amino acid sequence encoding an endoplasmic reticulum ("ER") retention domain that comprises an ER retention sequence. In one embodiment, the chimeric immunogen comprises the amino acid sequence of a non-toxic PE wherein domain Ib further comprises the non-native epitope between two cysteine residues of domain Ib.

In certain embodiments the cell recognition domain binds to $\alpha 2$ -macroglobulin receptor (" $\alpha 2$ -MR"), epidermal growth factor ("EGF") receptor, IL-2 receptor, IL-6 receptor, human transferrin receptor or gp120. In another embodiment, the cell recognition domain comprises amino acid sequences of a growth factor. In another embodiment, the translocation domain comprises amino acids 280 to 364 of domain II of PE. In another embodiment, the non-native epitope domain comprises a cysteine-cysteine loop that comprises the non-native epitope. In another embodiment, the non-native epitope domain comprises an amino acid sequence selected from the V3 loop

of HIV-1. In another embodiment, the ER retention domain is domain III of PE comprising a mutation that eliminates ADP ribosylation activity, such as Δ E553. The ER retention domain can comprise the ER retention sequence REDLK (SEQ ID NO:11), REDL (SEQ ID NO:12) or KDEL (SEQ ID NO:13). In another embodiment the non-native epitope is an epitope from a pathogen (e.g., an epitope from a virus, bacterium or parasitic protozoa) or from a cancer antigen.

In another embodiment the cell recognition domain is domain Ia of PE, the translocation domain is domain II of PE, the non-native epitope domain comprises an amino acid sequence encoding a non-native epitope inserted between two cysteine residues of domain Ib of PE, and the ER retention domain is domain III of PE and comprises a mutation that eliminates ADP ribosylation activity.

In another aspect, this invention provides a recombinant polynucleotide comprising a nucleotide sequence encoding a non-toxic *Pseudomonas* exotoxin A-like chimeric immunogen of this invention. In one embodiment, the recombinant polynucleotide is an expression vector further comprising an expression control sequence operatively linked to the nucleotide sequence.

In another aspect, this invention provides a recombinant *Pseudomonas* exotoxin A-like chimeric immunogen cloning platform comprising a nucleotide sequence encoding: (1) a cell recognition domain of between 10 and 1500 amino acids that binds to a cell surface receptor; (2) a translocation domain comprising an amino acid sequence substantially identical to a sequence of PE domain II sufficient to effect translocation to a cell cytosol; (3) an amino acid sequence encoding an endoplasmic reticulum ("ER") retention domain that comprises an ER retention sequence and, optionally, (4) a splicing site between the sequence encoding the translocation domain and the sequence encoding the ER retention domain. In one embodiment the recombinant polynucleotide is an expression vector further comprising an expression control sequence operatively linked to the nucleotide sequence.

In another aspect this invention provides a method of producing antibodies against a non-native epitope naturally within a cysteine-cysteine loop. The method comprises the step of inoculating an animal with a non-toxic *Pseudomonas* exotoxin A-like chimeric immunogen of this invention wherein the non-native epitope domain comprises a cysteine-cysteine loop that comprises the non-native epitope.

In another aspect this invention provides a vaccine comprising at least one *Pseudomonas* exotoxin A-like chimeric immunogen comprising a cell recognition domain, a translocation domain, a non-native epitope domain comprising a non-native epitope and an endoplasmic reticulum ("ER") retention domain comprising an ER retention sequence. In one embodiment the vaccine comprises a plurality of PE-like chimeric immunogens, each immunogen having a different non-native epitope. In another embodiment the different non-native epitopes are epitopes of different strains of the same pathogen.

In another aspect this invention provides a method of eliciting an immune response against a non-native epitope in a subject. The method comprises the step of administering to the subject a vaccine comprising at least one *Pseudomonas* exotoxin A-like chimeric immunogen of this invention. In one embodiment, the non-native epitope comprises a binding motif for an MHC Class II molecule of the subject and the immune response elicited is an MHC Class-II dependent cell-mediated immune response. In another embodiment the non-native epitope comprises a binding motif for an MHC Class I molecule of the subject and the immune response elicited is an MHC Class-I dependent cell-mediated immune response.

In another aspect this invention provides a polynucleotide vaccine comprising at least one recombinant polynucleotide comprising a nucleotide sequence encoding a non-toxic *Pseudomonas* exotoxin A-like chimeric immunogen of this invention.

In another aspect, this invention provides a method of eliciting an immune response against a non-native epitope in a subject. The method comprises the step of administering to the subject a polynucleotide vaccine comprising at least one recombinant polynucleotide comprising a nucleotide sequence encoding a non-toxic *Pseudomonas* exotoxin A-like chimeric immunogen of this invention. In one embodiment, the recombinant polynucleotide is an expression vector comprising an expression control sequence operatively linked to the nucleotide sequence.

In another aspect this invention provides a method of eliciting an immune response against a non-native epitope in a subject, the method comprising the steps of transfecting cells with a recombinant polynucleotide comprising a nucleotide sequence encoding a non-toxic *Pseudomonas* exotoxin A-like chimeric immunogen of this invention, and administering the cells to the subject.

In another aspect, this invention provides methods of eliciting an IgA-mediated secretory immune response. The methods involve administering to a mucosal membrane a non-toxic *Pseudomonas* chimeric immunogen of this invention, wherein the cell recognition domain binds to a receptor on a mucosal membrane. The cell
 5 recognition domain can bind to $\alpha 2$ -MR (e.g., the native cell recognition domain of PE), or to the EGF receptor. The mucosal surface can be mouth, nose, lung, gut, vagina, colon or rectum.

In another aspect, this invention provides a composition comprising secretory IgA antibodies that specifically recognize an epitope of a pathogen that enters
 10 the body through a mucosal surface, e.g., an epitope of HIV-1.

BRIEF DESCRIPTION OF THE DRAWINGS

Figs 1A-1C. (A and B) A schematic depiction of PE and a PE-V3 loop chimera showing the relative location of the Ib and V3 loops between domains II and III. Approximate location of the single amino acid deletion (Δ E553) to ablate PE toxicity is
 15 also shown. (C) Amino acid sequences, represented with single letter code, which replaced the Ib loop of wild-type PE with a V3 loop sequence of gp120 (bold type) from either the MN or Thai-E (TE) strains of HIV-1 contained two cysteine residues designed to result in a loop conformation following disulfide bond formation. The insertion of a
 20 unique PstI restriction site, used for introduction of V3 loop sequences, resulted in several modifications of the wild-type PE amino acid sequence adjacent to the Ib loop (italics). An irrelevant control peptide insert was prepared as a control and is designated ntPE-fp16. Calculated molecular masses are shown for full-length expressed proteins. Wild-type PE -- SEQ ID NO:6; ntPE-V3MN14 -- SEQ ID NO:7; ntPE-V3MN26 -- SEQ
 25 ID NO:8; ntPE-V3Th-E26 -- SEQ ID NO:9; ntPE-fp16 -- SEQ ID NO:10.

Figs 2A-2C. Characterization of ntPE-V3 loop chimeras after separation by SDS-PAGE. (A) Coomassie blue staining of purified ntPE-V3 loop chimeras following separation by SDS-PAGE. Approximately 1 μ g of protein was loaded on each lane. (B)
 30 Western blot analysis of ntPE-V3 loop chimeras. After transfer to Immobilon P membranes, proteins were probed with monoclonal antibodies raised against intact gp120/MN (1F12) or gp120/Thai-E (1B2). An irrelevant sequence of 16 amino adds was inserted into the Ib loop region of ntPE (ntPE-fp16) and was used here as a negative control. (C) Immunocapture studies, using either 1F12 or 1B2 immobilized on protein G

sepharose, were used to characterize the exposure of V3 loop sequences on the surface of the various chimeric proteins. Proteins were visualized by staining gels with Coomassie blue. Gp120 and ntPE-fp16 were used as positive and negative controls respectively. The capture of PE-V3 loop proteins is indicated with a single arrowhead and of gp120 by a double arrowhead. The left panel shows the presence of the antibody heavy chain (hc) only since the light chain (lc) was run off the gel. The right panel shows both chains.

Figs. 3A-3C. V3 loop amino acid sequence insertions do not significantly alter the secondary structure of wild-type PE. Near UV (A) and far UV (B) CD spectra (mean of three scans following background spectrum subtraction) were digitally smoothed, corrected for concentration, and normalized to units of mean residue weight ellipticity. (C) Secondary structure calculations were performed using the SELCON fitting program. *Calculated α -helix content agrees with values determined from changes in observed ellipticity at 222 nm.

Fig. 4. Toxic PE-V3 loop chimeras affect cell survival. The extent of protein synthesis, assessed by ^3H -leucine incorporation, was determined in human A431 cells following 18 h of exposure to various concentrations of either wild-type PE or a toxic form (with a glutamic acid residue at position 553 and capable of ADP ribosylating elongation factor 2) of PE-V3MN26.

Figs 5A-5B. Characterization of rabbit sera following immunization with either ntPE-V3MN26 or ntPE-V3Th-E26. (A) Western blot reactivity of rabbit antisera diluted 1:1000 for recombinant gp120/MN and gp120/Th-E was assessed following SDS-PAGE and the transfer of proteins to Immobilon P membranes. Reactive primary antibody was detected by a secondary anti-rabbit antibody conjugated to horseradish peroxidase. (B) Rabbit sera obtained from animals injected with ntPE-V3MN26 was pre-incubated with competing soluble gp120/MN at concentrations up to 50 $\mu\text{g}/\text{ml}$. Residual reactivity was detected by Western blot analysis of immobilized gp120/MN as described for (A).

Fig. 6. A ntPE-V3 loop chimera administered to rabbits produces an antibody response capable of neutralizing HIV-1 infectivity *in vitro*. Rabbits were immunized subcutaneously with 200 μg ntPE-V3MN26 and boosted similarly after 2, 4 and 12 weeks. Sera collected up to 27 weeks after the initial administration were evaluated for the ability to protect a human T-cell line, MT4, from killing by HIV-1 MN

as assessed by an MTT dye conversion assay. Values represent triplicate readings normalized against a control MT4 incubation not challenged by virus.

Fig. 7 is a diagram of *Pseudomonas* Exotoxin A structure. The amino acid position based on SEQ ID NO:2 is indicated. Domain 1a extends from amino acids 1-252. Domain II extends from amino acids 253-364. It includes a cysteine-cysteine loop formed by cysteines at amino acids 265-287. Furin cleaves within the cysteine-cysteine loop between amino acids 279 and 280. A fragment of PE beginning with amino acid 280 translocates to the cytosol. Constructs in which amino acids 345-364 are eliminated also translocate. Domain Ib spans amino acids 365-399. It contains a cysteine-cysteine loop formed by cysteines at amino acids 372 and 379. The domain can be eliminated entirely. Domain III spans amino acids 400-613. Deletion of amino acid 553 eliminates ADP ribosylation activity. The endoplasmic reticulum sequence, REDLK (SEQ ID NO:11) is located at the carboxy-terminus of the molecule, from amino acid 609-613.

Fig. 8 demonstrates that PE-V3 loop chimeras are trafficked similarly to native PE. Confluent monolayers of Caco-2 cells were exposed apically to recombinant, enzymatically-active *Pseudomonas* exotoxin (rEA-PE). Cell killing produced by 24 h of exposure at various native PE (rEA-PE) concentrations were compared to that produced by similar treatment with enzymatically-active versions of PE chimeras containing either 14 or 26 amino acids of the V3 loop of HIV-1 MNgp120.

Fig. 9 demonstrates that PE-V3 loop chimeras induce a serum IgG response. A non-toxic (enzymatically inactive) V3 loop chimera containing 26 amino acids of the V3 loop of HIV-1 MNgp120 (PEMN26) was administered to rabbits through six different inoculation protocols. Serum samples drawn at the times described were assayed by ELISA for MNgp120-specific IgG using a monoclonal antibody (1F12) which recognizes the V3 loop of this protein for assay calibration.

Fig. 10 shows that PE-V3 loop chimeras induce a salivary IgA response. A non-toxic (enzymatically inactive) V3 loop chimera containing 26 amino acids of the V3 loop of HIV-1 MNgp120 (PEMN26) was administered to rabbits through six different inoculation protocols. Saliva samples obtained following pilocarpine administration at the times described were assayed by ELISA for MNgp120-specific IgA. No gp120-specific IgA antibody was available for assay calibration. Values are reported as values normalized to a standardized positive sample.

Fig. 11 shows relative levels of salivary IgA following mucosal or systemic inoculation with ntPE-V3MN26. MN-gp120 specific IgA antibodies were measured by ELISA in saliva samples, normalized against a strongly positive sample and reported on an arbitrary scale of one antigen-specific IgA unit.

Fig. 12 shows serum levels of IgG following mucosal or systemic inoculation with ntPE-V3MN26. MN-gp120 specific IgG antibodies were measured in serum samples by ELISA and standardized against a mouse monoclonal antibody which specifically recognizes the V3 loop of MNgp120.

Fig. 13 shows serum levels of IgG following subcutaneous injection of ntPE-V3MN26. The immune response produced from injection of ntPE-V3MN26 (hatched bars) was compared to that induced when co-injected with a regimen of Freund's complete and incomplete adjuvant (solid bars). Non-toxic PE not containing the 26 amino acids from the V3 loop of MNgp120 was injected with the same adjuvant regimen as a control. MN-gp120 specific IgG antibodies were measured in serum samples by ELISA and standardized against a mouse monoclonal antibody which specifically recognizes the V3 loop of MNgp120.

Figs. 14A and 14B shows neutralization of clinical HIV isolates with antibodies elicited with the chimeric immunogens of this invention. Postvaccination sera from rabbits injected with ntPE-V3MN26 were mixed with either a B (Fig. 14A) or E (Fig. 14B) subtype virus. After a 1-h incubation at 37° C, viral infectivity was determined by adding treated virus to PBMCs for another 3 days. Inhibition of viral growth was evaluated by measuring p24 levels. Open square: p24 antigen (uninfected); closed circle: p24 antigen 1 prebleed sera; open circle: p24 antigen 1 immune sera (24 weeks).

DETAILED DESCRIPTION OF THE INVENTION

I. DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton et al., *DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY* (2d ed. 1994); *THE CAMBRIDGE DICTIONARY OF SCIENCE AND TECHNOLOGY* (Walker ed., 1988); *THE GLOSSARY OF*

GENETICS, 5TH ED., R. Rieger et al. (eds.), Springer Verlag (1991); and Hale & Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY (1991). As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

"Polynucleotide" refers to a polymer composed of nucleotide units.

5 Polynucleotides include naturally occurring nucleic acids, such as deoxyribonucleic acid ("DNA") and ribonucleic acid ("RNA") as well as nucleic acid analogs. Nucleic acid analogs include those which include non-naturally occurring bases, nucleotides that engage in linkages with other nucleotides other than the naturally occurring phosphodiester bond or which include bases attached through linkages other than phosphodiester bonds. Thus, nucleotide analogs include, for example and without
10 limitation, phosphorothioates, phosphorodithioates, phosphotriesters, phosphoramidates, boranophosphates, methylphosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs), and the like. Such polynucleotides can be synthesized, for example, using an automated DNA synthesizer. The term "nucleic acid" typically refers to large polynucleotides. The term "oligonucleotide" typically refers to
15 short polynucleotides, generally no greater than about 50 nucleotides. It will be understood that when a nucleotide sequence is represented by a DNA sequence (i.e., A, T, G, C), this also includes an RNA sequence (i.e., A, U, G, C) in which "U" replaces "T."

20 "cDNA" refers to a DNA that is complementary or identical to an mRNA, in either single stranded or double stranded form.

Conventional notation is used herein to describe polynucleotide sequences: the left-hand end of a single-stranded polynucleotide sequence is the 5'-end; the left-hand direction of a double-stranded polynucleotide sequence is referred to as the 5'-direction.
25 The direction of 5' to 3' addition of nucleotides to nascent RNA transcripts is referred to as the transcription direction. The DNA strand having the same sequence as an mRNA is referred to as the "coding strand"; sequences on the DNA strand having the same sequence as an mRNA transcribed from that DNA and which are located 5' to the 5'-end of the RNA transcript are referred to as "upstream sequences"; sequences on the DNA
30 strand having the same sequence as the RNA and which are 3' to the 3' end of the coding RNA transcript are referred to as "downstream sequences."

"Complementary" refers to the topological compatibility or matching together of interacting surfaces of two polynucleotides. Thus, the two molecules can be

described as complementary, and furthermore, the contact surface characteristics are complementary to each other. A first polynucleotide is complementary to a second polynucleotide if the nucleotide sequence of the first polynucleotide is identical to the nucleotide sequence of the polynucleotide binding partner of the second polynucleotide.

Thus, the polynucleotide whose sequence 5'-TATAC-3' is complementary to a polynucleotide whose sequence is 5'-GTATA-3'.

A nucleotide sequence is "substantially complementary" to a reference nucleotide sequence if the sequence complementary to the subject nucleotide sequence is substantially identical to the reference nucleotide sequence.

"Encoding" refers to the inherent property of specific sequences of nucleotides in a polynucleotide, such as a gene, a cDNA, or an mRNA, to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (i.e., rRNA, tRNA and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom. Thus, a gene encodes a protein if transcription and translation of mRNA produced by that gene produces the protein in a cell or other biological system. Both the coding strand, the nucleotide sequence of which is identical to the mRNA sequence and is usually provided in sequence listings, and non-coding strand, used as the template for transcription, of a gene or cDNA can be referred to as encoding the protein or other product of that gene or cDNA. Unless otherwise specified, a "nucleotide sequence encoding an amino acid sequence" includes all nucleotide sequences that are degenerate versions of each other and that encode the same amino acid sequence. Nucleotide sequences that encode proteins and RNA may include introns.

"Recombinant polynucleotide" refers to a polynucleotide having sequences that are not naturally joined together. An amplified or assembled recombinant polynucleotide may be included in a suitable vector, and the vector can be used to transform a suitable host cell. A host cell that comprises the recombinant polynucleotide is referred to as a "recombinant host cell." The gene is then expressed in the recombinant host cell to produce, e.g., a "recombinant polypeptide." A recombinant polynucleotide may serve a non-coding function (e.g., promoter, origin of replication, ribosome-binding site, etc.) as well.

"Expression control sequence" refers to a nucleotide sequence in a polynucleotide that regulates the expression (transcription and/or translation) of a

nucleotide sequence operatively linked thereto. "Operatively linked" refers to a functional relationship between two parts in which the activity of one part (e.g., the ability to regulate transcription) results in an action on the other part (e.g., transcription of the sequence). Expression control sequences can include, for example and without
5 limitation, sequences of promoters (e.g., inducible or constitutive), enhancers, transcription terminators, a start codon (i.e., ATG), splicing signals for introns, and stop codons.

"Expression vector" refers to a vector comprising a recombinant polynucleotide comprising expression control sequences operatively linked to a nucleotide
10 sequence to be expressed. An expression vector comprises sufficient *cis*-acting elements for expression; other elements for expression can be supplied by the host cell or *in vitro* expression system. Expression vectors include all those known in the art, such as cosmids, plasmids (e.g., naked or contained in liposomes) and viruses that incorporate the recombinant polynucleotide.

"Amplification" refers to any means by which a polynucleotide sequence is copied and thus expanded into a larger number of polynucleotide molecules, e.g., by
15 reverse transcription, polymerase chain reaction, and ligase chain reaction.

"Primer" refers to a polynucleotide that is capable of specifically hybridizing to a designated polynucleotide template and providing a point of initiation for
20 synthesis of a complementary polynucleotide. Such synthesis occurs when the polynucleotide primer is placed under conditions in which synthesis is induced, i.e., in the presence of nucleotides, a complementary polynucleotide template, and an agent for polymerization such as DNA polymerase. A primer is typically single-stranded, but may be double-stranded. Primers are typically deoxyribonucleic acids, but a wide variety of
25 synthetic and naturally occurring primers are useful for many applications. A primer is complementary to the template to which it is designed to hybridize to serve as a site for the initiation of synthesis, but need not reflect the exact sequence of the template. In such a case, specific hybridization of the primer to the template depends on the stringency of the hybridization conditions. Primers can be labeled with, e.g.,
30 chromogenic, radioactive, or fluorescent moieties and used as detectable moieties.

"Probe," when used in reference to a polynucleotide, refers to a polynucleotide that is capable of specifically hybridizing to a designated sequence of another polynucleotide. A probe specifically hybridizes to a target complementary

polynucleotide, but need not reflect the exact complementary sequence of the template. In such a case, specific hybridization of the probe to the target depends on the stringency of the hybridization conditions. Probes can be labeled with, e.g., chromogenic, radioactive, or fluorescent moieties and used as detectable moieties.

5 A first sequence is an "antisense sequence" with respect to a second sequence if a polynucleotide whose sequence is the first sequence specifically hybridizes with a polynucleotide whose sequence is the second sequence.

"Hybridizing specifically to" or "specific hybridization" or "selectively hybridize to", refers to the binding, duplexing, or hybridizing of a nucleic acid molecule preferentially to a particular nucleotide sequence under stringent conditions when that sequence is present in a complex mixture (e.g., total cellular) DNA or RNA.

10 The term "stringent conditions" refers to conditions under which a probe will hybridize preferentially to its target subsequence, and to a lesser extent to, or not at all to, other sequences. "Stringent hybridization" and "stringent hybridization wash conditions" in the context of nucleic acid hybridization experiments such as Southern and northern hybridizations are sequence dependent, and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes* part I chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays", Elsevier, New York. Generally, highly stringent hybridization and wash conditions are selected to be about 5° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very 20 stringent conditions are selected to be equal to the T_m for a particular probe.

25 An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is 50% formalin with 1 mg of heparin at 42° C, with the hybridization being carried out overnight. An example of highly stringent wash conditions is 0.15 M NaCl at 72° C for about 15 minutes. An example of stringent wash 30 conditions is a 0.2X SSC wash at 65° C for 15 minutes (see, Sambrook et al. for a description of SSC buffer). Often, a high stringency wash is preceded by a low stringency wash to remove background probe signal. An example medium stringency

wash for a duplex of, e.g., more than 100 nucleotides, is 1x SSC at 45° C for 15 minutes. An example low stringency wash for a duplex of, e.g., more than 100 nucleotides, is 4-6x SSC at 40° C for 15 minutes. In general, a signal to noise ratio of 2x (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization.

"Polypeptide" refers to a polymer composed of amino acid residues, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof linked via peptide bonds, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof. Synthetic polypeptides can be synthesized, for example, using an automated polypeptide synthesizer. The term "protein" typically refers to large polypeptides. The term "peptide" typically refers to short polypeptides.

Conventional notation is used herein to portray polypeptide sequences: the left-hand end of a polypeptide sequence is the amino-terminus; the right-hand end of a polypeptide sequence is the carboxyl-terminus.

"Conservative substitution" refers to the substitution in a polypeptide of an amino acid with a functionally similar amino acid. The following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

"Allelic variant" refers to any of two or more polymorphic forms of a gene occupying the same genetic locus. Allelic variations arise naturally through mutation, and may result in phenotypic polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. "Allelic variants" also refer to cDNAs derived from mRNA transcripts of genetic allelic variants, as well as the proteins encoded by them.

The terms "identical" or percent "identity," in the context of two or more polynucleotide or polypeptide sequences, refer to two or more sequences or subsequences

that are the same or have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection.

5 The phrase "substantially identical," in the context of two nucleic acids or polypeptides, refers to two or more sequences or subsequences that have at least 60%, 80%, 90%, 95% or 98% nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection. Preferably, the substantial identity exists over a region of the sequences that is at least about 50 residues in length, more preferably over a region of at least about 100 residues, and most preferably the sequences are substantially identical over at least about 150 residues. In a most preferred embodiment, the sequences are substantially identical over the entire length of the coding regions.

15 For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

25 Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (*see generally* Ausubel *et al.*, *supra*).

30 One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP

uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35:351-360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. For example, a reference sequence can be compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps.

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or

the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP
5 program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)).

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g.,
10 Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1,
15 more preferably less than about 0.01, and most preferably less than about 0.001.

A further indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid,
20 as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions, as described herein.

25 A "ligand" is a compound that specifically binds to a target molecule.

A "receptor" is compound that specifically binds to a ligand.

"Antibody" refers to a polypeptide ligand substantially encoded by an immunoglobulin gene or immunoglobulin genes, or fragments thereof, which specifically binds and recognizes an epitope (e.g., an antigen). The recognized immunoglobulin
30 genes include the kappa and lambda light chain constant region genes, the alpha, gamma, delta, epsilon and mu heavy chain constant region genes, and the myriad immunoglobulin variable region genes. Antibodies exist, e.g., as intact immunoglobulins or as a number of well characterized fragments produced by digestion with various peptidases. This

includes, e.g., Fab' and F(ab')₂ fragments. The term "antibody," as used herein, also includes antibody fragments either produced by the modification of whole antibodies or those synthesized *de novo* using recombinant DNA methodologies. It also includes polyclonal antibodies, monoclonal antibodies, chimeric antibodies and humanized antibodies. "Fc" portion of an antibody refers to that portion of an immunoglobulin heavy chain that comprises one or more heavy chain constant region domains, CH₁, CH₂ and CH₃, but does not include the heavy chain variable region.

A ligand or a receptor (e.g., an antibody) "specifically binds to" or "is specifically immunoreactive with" a compound analyte when the ligand or receptor functions in a binding reaction which is determinative of the presence of the analyte in a sample of heterogeneous compounds. Thus, under designated assay (e.g., immunoassay) conditions, the ligand or receptor binds preferentially to a particular analyte and does not bind in a significant amount to other compounds present in the sample. For example, a polynucleotide specifically binds under hybridization conditions to an analyte polynucleotide comprising a complementary sequence; an antibody specifically binds under immunoassay conditions to an antigen analyte bearing an epitope against which the antibody was raised; and an adsorbent specifically binds to an analyte under proper elution conditions.

"Immunoassay" refers to a method of detecting an analyte in a sample involving contacting the sample with an antibody that specifically binds to the analyte and detecting binding between the antibody and the analyte. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York, for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity.

"Vaccine" refers to an agent or composition containing an agent effective to confer a therapeutic degree of immunity on an organism while causing only very low levels of morbidity or mortality. Methods of making vaccines are, of course, useful in the study of the immune system and in preventing and treating animal or human disease.

An "immunogenic amount" is an amount effective to elicit an immune response in a subject.

"Substantially pure" or "isolated" means an object species is the predominant species present (i.e., on a molar basis, more abundant than any other individual macromolecular species in the composition), and a substantially purified fraction is a composition wherein the object species comprises at least about 50% (on a molar basis) of all macromolecular species present. Generally, a substantially pure composition means that about 80% to 90% or more of the macromolecular species present in the composition is the purified species of interest. The object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) if the composition consists essentially of a single macromolecular species. Solvent species, small molecules (< 500 Daltons), stabilizers (e.g., BSA), and elemental ion species are not considered macromolecular species for purposes of this definition.

"Naturally-occurring" as applied to an object refers to the fact that the object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring.

"Detecting" refers to determining the presence, absence, or amount of an analyte in a sample, and can include quantifying the amount of the analyte in a sample or per cell in a sample.

"Detectable moiety" or a "label" refers to a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include ^{32}P , ^{35}S , fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin-streptavidin, dioxigenin, haptens and proteins for which antisera or monoclonal antibodies are available, or nucleic acid molecules with a sequence complementary to a target. The detectable moiety often generates a measurable signal, such as a radioactive, chromogenic, or fluorescent signal, that can be used to quantitate the amount of bound detectable moiety in a sample. The detectable moiety can be incorporated in or attached to a primer or probe either covalently, or through ionic, van der Waals or hydrogen bonds, e.g., incorporation of radioactive nucleotides, or biotinylated nucleotides that are recognized by streptavidin. The detectable moiety may be directly or indirectly detectable. Indirect detection can involve the binding of a second directly or indirectly detectable moiety to the detectable

moiety. For example, the detectable moiety can be the ligand of a binding partner, such as biotin, which is a binding partner for streptavidin, or a nucleotide sequence, which is the binding partner for a complementary sequence, to which it can specifically hybridize. The binding partner may itself be directly detectable, for example, an antibody may be itself labeled with a fluorescent molecule. The binding partner also may be indirectly detectable, for example, a nucleic acid having a complementary nucleotide sequence can be a part of a branched DNA molecule that is in turn detectable through hybridization with other labeled nucleic acid molecules. (See, e.g., PD. Fahrlander and A. Klausner, *Bio/Technology* (1988) 6:1165.) Quantitation of the signal is achieved by, e.g., scintillation counting, densitometry, or flow cytometry.

"Linker" refers to a molecule that joins two other molecules, either covalently, or through ionic, van der Waals or hydrogen bonds, e.g., a nucleic acid molecule that hybridizes to one complementary sequence at the 5' end and to another complementary sequence at the 3' end, thus joining two non-complementary sequences.

"Pharmaceutical composition" refers to a composition suitable for pharmaceutical use in a mammal. A pharmaceutical composition comprises a pharmacologically effective amount of an active agent and a pharmaceutically acceptable carrier. "Pharmacologically effective amount" refers to that amount of an agent effective to produce the intended pharmacological result. "Pharmaceutically acceptable carrier" refers to any of the standard pharmaceutical carriers, buffers, and excipients, such as a phosphate buffered saline solution, 5% aqueous solution of dextrose, and emulsions, such as an oil/water or water/oil emulsion, and various types of wetting agents and/or adjuvants. Suitable pharmaceutical carriers and formulations are described in *Remington's Pharmaceutical Sciences*, 19th Ed. (Mack Publishing Co., Easton, 1995). Preferred pharmaceutical carriers depend upon the intended mode of administration of the active agent. Typical modes of administration include enteral (e.g., oral) or parenteral (e.g., subcutaneous, intramuscular, intravenous or intraperitoneal injection; or topical, transdermal, or transmucosal administration). A "pharmaceutically acceptable salt" is a salt that can be formulated into a compound for pharmaceutical use including, e.g., metal salts (sodium, potassium, magnesium, calcium, etc.) and salts of ammonia or organic amines.

"Small organic molecule" refers to organic molecules of a size comparable to those organic molecules generally used in pharmaceuticals. The term excludes organic

biopolymers (e.g., proteins, nucleic acids, etc.). Preferred small organic molecules range in size up to about 5000 Da, up to about 2000 Da, or up to about 1000 Da.

A "subject" of diagnosis or treatment is a human or non-human animal, including a mammal or a primate.

5 "Treatment" refers to prophylactic treatment or therapeutic treatment.

A "prophylactic" treatment is a treatment administered to a subject who does not exhibit signs of a disease or exhibits only early signs for the purpose of decreasing the risk of developing pathology.

A "therapeutic" treatment is a treatment administered to a subject who exhibits signs of pathology for the purpose of diminishing or eliminating those signs.

"Diagnostic" means identifying the presence or nature of a pathologic condition. Diagnostic methods differ in their specificity and selectivity. While a particular diagnostic method may not provide a definitive diagnosis of a condition, it suffices if the method provides a positive indication that aids in diagnosis.

"Prognostic" means predicting the probable development (e.g., severity) of a pathologic condition.

"Plurality" means at least two.

"*Pseudomonas* exotoxin A" or "PE" is secreted by *Ps aeruginosa* as a 67 kD protein composed of three prominent globular domains (Ia, II, and III) and one small subdomain (Ib) connecting domains II and III. (A.S. Allured et. al. (1986) *Proc. Natl. Acad. Sci.* 83:1320-1324.) Domain Ia of PE mediates cell binding. In nature, domain Ia binds to the low density lipoprotein receptor-related protein ("LRP"), also known as the α 2-macroglobulin receptor (" α 2-MR"). (M.Z. Kounnas *et al.* (1992) *J. Biol. Chem.* 267:12420-23.) It spans amino acids 1-252. Domain II mediates translocation to the cytosol. It spans amino acids 253-364. Domain Ib has no known function. It spans amino acids 365-399. Domain III is responsible for cytotoxicity and includes an endoplasmic reticulum retention sequence. It mediates ADP ribosylation of elongation factor 2, which inactivates protein synthesis. It spans amino acids 400-613. PE is "non-toxic" if it lacks EF2 ADP ribosylation activity. Deleting amino acid E553 (" Δ E553") from domain III detoxifies the molecule. PE having the mutation Δ E553 is referred to herein as "PE Δ E553." Genetically modified forms of PE are described in, e.g., Pastan *et al.*, United States patent 5,602,095; Pastan *et al.*, United States patent 5,512,658 and Pastan *et al.*, United States patent 5,458,878. Allelic forms of PE are included in this

definition. See, e.g., M.L. Vasil *et al.*, (1986) *Infect. Immunol.* 52:538-48.

"Cysteine-cysteine loop" refers to a peptide moiety in a polypeptide that is defined by an amino acid sequence bordered by two disulfide-bonded cysteine residues.

"Non-native epitope" refers to an epitope encoded by an amino acid sequence that does not naturally occur in the Ib domain of *Pseudomonas* exotoxin A.

II. PSEUDOMONAS EXOTOXIN A-LIKE CHIMERIC IMMUNOGENS

A. Basic Structure

The *Pseudomonas* exotoxin A-like ("PE-like") chimeric immunogens of this invention are polypeptides having structural domains organized, except as provided herein, in the same sequence as the four structural domains of PE (i.e., Ia, II, Ib and III), and having certain functions (e.g., cell recognition, cytosolic translocation and endoplasmic reticulum retention) also possessed by the functional domains of PE. Additionally, the PE-like chimeric immunogens of this invention possess a domain that functionalizes a domain of PE for which no function yet has been identified. Namely, PE-like chimeric immunogens replace the Ib domain of PE with a functional non-native epitope domain that serves as an immunogen to elicit an immune response against the non-native epitope.

Accordingly, PE-like chimeric immunogens include the following structural domains comprised of amino acid sequences, the domains imparting particular functions to the chimeric protein: (1) a "cell recognition domain" that functions as a ligand for a cell surface receptor and that mediates binding of the protein to a cell; (2) a "translocation domain" that mediates translocation from the endosomes to the cytosol; (3) a "non-native epitope domain" that contains the immunogenic non-native epitope; and, optionally, (4) an "endoplasmic reticulum ("ER") retention domain" that functions to translocate the molecule from the endosome to the endoplasmic reticulum, from which it enters the cytosol. When the ER retention domain is eliminated the chimeric immunogen still can retain immunogenic function.

In one embodiment, a PE-like chimeric immunogen comprises the native sequence of PE, except for the Ib domain, which is engineered to include the amino acid sequence of a non-native epitope. For example, one can insert an amino acid sequence encoding the non-native epitope into the cysteine-cysteine loop of the Ib domain.

However, the relationship of PE structure to its function has been extensively studied. The amino acid sequence of PE has been re-engineered to provide new functions, and many amino acids or peptide segments critical and non-critical to PE function have been identified. The PE-like chimeric immunogens of this invention can incorporate these structural modifications to PE.

B. Cell Recognition Domain

The *Pseudomonas* exotoxin chimeras of this invention comprise an amino acid sequence encoding a "cell recognition domain." The cell recognition domain functions as a ligand for a cell surface receptor. It mediates binding of the protein to a cell. Its purpose is to target the chimera to a cell which will transport it to the cytosol for processing. The cell recognition domain can be located in the position of domain Ia of PE. However, this domain can be moved out of the normal organizational sequence. More particularly, the cell recognition domain can be inserted upstream of the ER retention domain. Alternatively the cell recognition domain can be chemically coupled to the toxin. Also, the chimera can include a first cell recognition domain at the location of the Ia domain and a second cell recognition domain upstream of the ER retention domain. Such constructs can bind to more than one cell type. See, e.g., R.J. Kreitman *et al.* (1992) *Bioconjugate Chem.* 3:63-68.

Because the cell recognition domain functions as a handle to attach the chimera to a cell, it can have the structure of any polypeptide known to bind to a particular receptor. Accordingly, the domain generally has the size of known polypeptide ligands, e.g., between about 10 amino acids and about 1500 amino acids, or about 100 amino acids and about 300 amino acids.

Several methods are useful for identifying functional cell recognition domains for use in chimeric immunogens. One method involves detecting binding between a chimera that comprises the cell recognition domain with the receptor or with a cell bearing the receptor. Other methods involve detecting entry of the chimera into the cytosol, indicating that the first step, cell binding, was successful. These methods are described in detail below in the section on testing.

The cell recognition domain can have the structure of any polypeptide that binds to a cell surface receptor. In one embodiment, the amino acid sequence is that of domain Ia of PE, thereby targeting the chimeric protein to the $\alpha 2$ -MR domain. In other

embodiments domain Ia can be substituted with: growth factors, such as TGF α , which binds to epidermal growth factor ("EGF"); IL-2, which binds to the IL-2 receptor; IL-6, which binds to the IL-6 receptor (e.g., activated B cells and liver cells); CD4, which binds to HIV-infected cells); a chemokine (e.g., Rantes, MIP-1 α or MIP-1 β), which binds to a chemokine receptor (e.g., CCR5 or fusin (CXCR4)); ligands for leukocyte cell surface receptors, for example, GM-CSF, G-CSF; ligands for the IgA receptor; or antibodies or antibody fragments directed to any receptor (e.g., single chain antibodies against human transferrin receptor). I. Pastan *et al.* (1992) *Annu. Rev. Biochem.* 61:331-54.

In one embodiment, the cell recognition domain is located in place of domain Ia of PE. It can be attached to the other moiety of the molecule through a linker. However, engineering studies show that *Pseudomonas* exotoxin can be targeted to certain cell types by introducing a cell recognition domain upstream of the ER retention sequence, which is located at the carboxy-terminus of the polypeptide. For example, TGF α has been inserted into domain III just before amino acid 604, i.e., about ten amino acids from the carboxy-terminus. This chimeric protein binds to cells bearing EGF receptor. Pastan *et al.*, U.S. Patent 5,602,095.

Cell specific ligands which are proteins can often be formed in part or in whole as a fusion protein with the *Pseudomonas* exotoxin chimeras of the present invention. A "fusion protein" refers to a polypeptide formed by the joining of two or more polypeptides through a peptide bond formed by the amino terminus of one polypeptide and the carboxyl terminus of the other polypeptide. The fusion protein may be formed by the chemical coupling of the constituent polypeptides but is typically expressed as a single polypeptide from a nucleic acid sequence encoding the single contiguous fusion protein. Included among such fusion proteins are single chain Fv fragments (scFv). Particularly preferred targeted *Pseudomonas* exotoxin chimeras are disulfide stabilized proteins which can be formed in part as a fusion protein as exemplified herein. Other protein cell specific ligands can be formed as fusion proteins using cloning methodologies well known to the skilled artisan.

Attachment of cell specific ligands also can be accomplished through the use of linkers. The linker is capable of forming covalent bonds or high-affinity non-covalent bonds to both molecules. Suitable linkers are well known to those of ordinary skill in the art and include, but are not limited to, straight or branched-chain

carbon linkers, heterocyclic carbon linkers, or peptide linkers. The linkers may be joined to the constituent amino acids through their side groups (*e.g.*, through a disulfide linkage to cysteine).

In one embodiment, domain Ia is replaced with a polypeptide sequence for an immunoglobulin heavy chain from an immunoglobulin specific for the target cell. The light chain of the immunoglobulin can be co-expressed with the PE-like chimeric immunogen so as to form a light chain-heavy chain dimer. In the conjugate protein, the antibody is chemically linked to a polypeptide comprising the other domains of the chimeric immunogen.

The procedure for attaching a *Pseudomonas* exotoxin chimera to an antibody or other cell specific ligand will vary according to the chemical structure of the toxin. Antibodies contain a variety of functional groups; *e.g.*, sulfhydryl (-S), carboxylic acid (COOH) or free amine (-NH₂) groups, which are available for reaction with a suitable functional group on a toxin. Additionally, or alternatively, the antibody or *Pseudomonas* exotoxin chimera can be derivatized to expose or attach additional reactive functional groups. The derivatization may involve attachment of any of a number of linker molecules such as those available from Pierce Chemical Company, Rockford Illinois.

A bifunctional linker having one functional group reactive with a group on the *Pseudomonas* exotoxin chimera, and another group reactive with a cell specific ligand, can be used to form a desired conjugate. Alternatively, derivatization may involve chemical treatment of the *Pseudomonas* exotoxin chimera or the cell specific ligand, *e.g.*, glycol cleavage of the sugar moiety of a glycoprotein antibody with periodate to generate free aldehyde groups. The free aldehyde groups on the antibody may be reacted with free amine or hydrazine groups on the antibody to bind the *Pseudomonas* exotoxin chimera thereto. (See J.D. Rodwell et al., U.S. Patent No. 4,671,958.) Procedures for generation of free sulfhydryl groups on antibodies or other proteins, are also known. (See R.A. Nicoletti et al., U.S. Pat. No. 4,659,839.)

C. Translocation Domain

PE-like chimeric immunogens also comprise an amino acid sequence encoding a "PE translocation domain." The PE translocation domain comprises an amino acid sequence sufficient to effect translocation of chimeric proteins that have been

endocytosed by the cell into the cytosol. The amino acid sequence is identical to, or substantially identical to, a sequence selected from domain II of PE.

The amino acid sequence sufficient to effect translocation can derive from the translocation domain of native PE. This domain spans amino acids 253-364. The translocation domain can include the entire sequence of domain II. However, the entire sequence is not necessary for translocation. For example, the amino acid sequence can minimally contain, e.g., amino acids 280-344 of domain II of PE. Sequences outside this region, i.e., amino acids 253-279 and/or 345-364, can be eliminated from the domain. This domain also can be engineered with substitutions so long as translocation activity is retained.

The translocation domain functions as follows. After binding to a receptor on the cell surface, the chimeric proteins enter the cell by endocytosis through clathrin-coated pits. Residues 265 and 287 are cysteines that form a disulfide loop. Once internalized into endosomes having an acidic environment, the peptide is cleaved by the protease furin between Arg279 and Gly280. Then, the disulfide bond is reduced. A mutation at Arg279 inhibits proteolytic cleavage and subsequent translocation to the cytosol. M. Ogata *et al.* (1990) *J. Biol. Chem.* 265:20678-85. However, a fragment of PE containing the sequence downstream of Arg279 (called "PE37") retains substantial ability to translocate to the cytosol. C.B. Siegall *et al.* (1989) *J. Biol. Chem.* 264:14256-61. Sequences in domain II beyond amino acid 345 also can be deleted without inhibiting translocation. Furthermore, amino acids at positions 339 and 343 appear to be necessary for translocation. C.B. Siegall *et al.* (1991) *Biochemistry* 30:7154-59.

Methods for determining the functionality of a translocation domain are described below in the section on testing.

D. Non-native Epitope Domain

PE-like chimeric immunogens also comprise an amino acid sequence encoding a "non-native epitope domain." The non-native epitope domain comprises the amino acid sequence of a non-native epitope. The domain functions to contain the immunogenic non-native epitope for presentation to the immune system. The non-native epitope domain is engineered into the Ib domain location of PE, between the translocation domain (e.g., domain II) and the ER retention domain (e.g., domain III).

Methods of determining immunogenicity of a translocation domain are described below in the section on testing.

The non-native epitope can be any amino acid sequence that is immunogenic. The non-native epitope domain can have between about 5 amino acids and about 1500 amino acids. This includes domains having between about 15 amino acids and about 350 amino acids or about 15 amino acids and about 50 amino acids.

In native *Pseudomonas* exotoxin A, domain Ib spans amino acids 365 to 399. The native Ib domain is structurally characterized by a disulfide bond between two cysteines at positions 372 and 379. Domain Ib is not essential for cell binding, translocation, ER retention or ADP ribosylation activity. Therefore, it can be entirely re-engineered.

The non-native epitope domain can be linear or it can include a cysteine-cysteine loop that comprises the non-native epitope. In one embodiment, the non-native epitope domain includes a cysteine-cysteine loop that comprises the non-native epitope. This arrangement offers several advantages. First, when the non-native epitope naturally exists inside, or comprises, a cysteine-cysteine disulfide bonded loop, the non-native epitope domain will present the epitope in near-native conformation. Second, it is believed that charged amino acid residues in the native Ib domain result in a hydrophilic structure that sticks out away from the molecule and into the solvent, where it is available to interact with immune system components. Therefore, placing the non-native epitope within a cysteine-cysteine loop results in more effective presentation when the non-native epitope also is hydrophilic. Third, the Ib domain is highly insensitive to mutation. Therefore, although the cysteine-cysteine loop of the native Ib domain has only six amino acids between the cysteine residues, one can insert much longer sequences into the loop without disrupting cell binding, translocation, ER retention or ADP ribosylation activity.

This invention envisions several ways in which to engineer the non-native epitope domain into the Ib domain location. One method involves inserting the amino acid sequence of the non-native epitope directly into the amino acid sequence of the Ib domain, with or without deletion of native amino acid sequences. Another method involves removing all or part of the Ib domain and replacing it with an amino acid sequence that includes the non-native epitope between two cysteine residues so that the cysteines engage in a disulfide bond when the protein is expressed. For example, if the

non-native epitope normally exists within a cysteine-cysteine loop structure of a polypeptide, a portion of the polypeptide that includes the loop and the non-native epitope can be inserted in place of the cysteine-cysteine loop domain.

The choice of the non-native epitope is at the discretion of the practitioner.

5 In choosing, the practitioner may consider the following. While the non-native epitope domain can be linear, non-native epitopes that naturally exist within a cysteine-cysteine loop take advantage of the natural structure of the Ib loop of *Pseudomonas* exotoxin A. Epitopes from agents responsible for indolent infections or cancer-specific antigens are attractive because these antigens tend to resist attack from the immune system. Also, recombinant technology allows one to quickly insert a polynucleotide encoding an epitope
10 into a vector encoding the chimeric protein. Therefore, one can quickly change sequences as a non-native epitope changes. Accordingly, epitopes from rapidly evolving infectious agents make attractive inserts.

Thus, for example, epitopes can be chosen from any pathogen, e.g.,
15 viruses, bacteria and protozoan parasites. Viral sources of epitopes include, for example, HIV, herpes zoster, influenza, polio and hepatitis. Bacterial sources include, for example, tuberculosis, Chlamydia or Salmonella. Parasitic protozoan sources include, for example, *Trypanosoma* or *Plasmodium*. In particular, the agent can be one that gains entry into the body through epithelial mucosal membranes. Useful cancer-specific antigens include those that are expressed on the cell surface and, therefore, can
20 be target of a cytotoxic T-lymphocyte response, such as a prostate cancer-specific marker (e.g., PSA), a breast cancer-specific marker (e.g., BRCA-1 or HER2), a pancreatic cancer-specific marker (e.g., CA9-19), a melanoma marker (e.g., tyrosinase) or a cancer-specific mutant form of EGF.

25 In one embodiment, the non-native epitope derives from the principal neutralizing loop of a retrovirus, such as HIV-1 or HIV-2. In particular, the epitope can derive from the V3 loop of gp120 protein from HIV-1. A neutralizing loop can be identified by neutralizing antibodies, i.e., antibodies that neutralize infectivity of the virus. The sequences can be from any strain, in particular, circulating strains. Such
30 strains include, for example, MN (e.g., subtype B) or Thai-E (e.g., subtype E). V3 loops of various strains of HIV-1 have about 35 amino acids. The strains of HIV can be T-cell tropic or macrophage tropic. In one embodiment, the sequences from the V3 loop include at least 8 amino acids (e.g., a peptide sufficiently long to fit into an MHC Class

II binding pocket) that includes a V3 loop apex. The V3 loop of MN strain of HIV has the sequence: CTRPNYNKRK RIHIGPGRAF YTTKNIIGTI RQAHC (SEQ ID NO:3). The V3 loop of Thai-E strain of HIV has the sequence: CTRPSNNTRT SITIGPGQVF YRTGDIIGDI RKAYC (SEQ ID NO:4). The V3 loop apex is underlined. The sequence be around 14 to around 26 amino acids long. A vaccine can comprise a plurality of immunogens having different viral epitopes.

In another embodiment the non-native epitope can be an epitope expressed by a cell during disease. For example, the non-native epitope can be a cancer cell marker. For example, certain breast cancers express a mutant EGF ("epidermal growth factor") receptor that results from a splice variant. This mutant form exhibits a unique epitope.

E. ER Retention Domain

PE-like chimeric immunogens also can comprise an amino acid sequence encoding an "endoplasmic reticulum retention domain." The endoplasmic reticulum ("ER") retention domain functions in translocating the chimeric protein to from the endosome to the endoplasmic reticulum, from where it is transported to the cytosol. The ER retention domain is located at the position of domain III in PE. The ER retention domain comprises an amino acid sequence that has, at its carboxy terminus, an ER retention sequence. The ER retention sequence in native PE is REDLK (SEQ ID NO:11). Lysine can be eliminated (i.e., REDL (SEQ ID NO:12)) without a decrease in activity. REDLK (from SEQ ID NO:1) can be replaced with other ER retention sequences, such as KDEL (SEQ ID NO:12), or polymers of these sequences. M. Ogata *et al.* (1990) *J. Biol. Chem.* 265:20678-85. Pastan *et al.*, U.S. Patent 5,458,878. I. Pastan *et al.* (1992) *Annu. Rev. Biochem.* 61:331-54.

Sequences up-stream of the ER retention sequence can be the native PE domain III (preferably de-toxified), can be entirely eliminated, or can be replaced by another amino acid sequence. If replaced by another amino acid sequence, the sequence can, itself, be highly immunogenic or can be slightly immunogenic. A highly immunogenic ER retention domain is preferable for use in eliciting a humoral immune response. Chimeras in which the ER retention domain is only slightly immunogenic will be more useful when an MHC Class I-dependent cell-mediated immune response is desired.

Activity of this domain can be assessed by testing for translocation of the protein into the target cell cytosol using the assays described below.

In native PE, the ER retention sequence is located at the carboxy terminus of domain III. Domain III has two functions in PE. It exhibits ADP-ribosylating activity and directs endocytosed toxin into the endoplasmic reticulum. Eliminating the ER retention sequence from the chimeric protein does not alter the activity of *Pseudomonas* exotoxin as a superantigen, but does inhibit its utility to elicit an MHC Class I-dependent cell-mediated immune response.

The ribosylating activity of PE is located between about amino acids 400 and 600 of PE. In methods of vaccinating a subject using the chimeric proteins of this invention, it is preferable that the protein be non-toxic. One method of doing so is by eliminating ADP ribosylation activity. In this way, the chimeric protein can function as a vector for non-native epitope sequences to be processed by the cell and presented on the cell surface with MHC Class I molecules, rather than as a toxin. ADP ribosylation activity can be eliminated by, for example, deleting amino acid E553 ("ΔE553"). M. Lukac *et al.* (1988) *Infect. and Immun.* 56:3095-3098. Alternatively, the amino acid sequence of domain III, or portions of it, can be deleted from the protein. Of course, an ER retention sequence should be included at the carboxy-terminus.

In one embodiment, the sequence of the ER retention domain is substantially identical to the native amino acid sequences of the domain III, or a fragment of it. In one embodiment, the ER retention domain is domain III of PE.

In another embodiment, a cell recognition domain is inserted into the amino acid sequence of the ER retention domain (e.g., into domain III). For example, the cell recognition domain can be inserted just up-stream of the ER retention sequence, so that the ER retention sequence is connected directly or within ten amino acids of the carboxy terminus of the cell recognition domain.

F. Methods Of Making PE-like Chimeric Immunogens

PE-like chimeric immunogens preferably are produced recombinantly, as described below. This invention also envisions the production of PE chimeric proteins by chemical synthesis using methods available to the art.

G. Testing PE-like Immunogenic Chimeras

Having selected various structures as domains of the chimeric immunogen, the function of these domains, and of the chimera as a whole, can be tested to detect functionality. PE-like immunogenic chimeras can be tested for cell recognition, cytosolic translocation and immunogenicity using routine assays. The entire chimeric protein can be tested, or, the function of various domains can be tested by substituting them for native domains of the wild-type toxin.

1. **Receptor binding/Cell recognition**

The function of the cell binding domain can be tested as a function of the chimera to bind to the target receptor either isolated or on the cell surface.

In one method, binding of the chimera to a target is performed by affinity chromatography. For example, the chimera can be attached to a matrix in an affinity column, and binding of the receptor to the matrix detected.

Binding of the chimera to receptors on cells can be tested by, for example, labeling the chimera and detecting its binding to cells by, e.g., fluorescent cell sorting, autoradiography, etc.

If antibodies have been identified that bind to the ligand from which the cell recognition domain is derived, they also are useful to detect the existence of the cell recognition domain in the chimeric immunogen by immunoassay, or by competition assay for the cognate receptor.

2. **Translocation to the cytosol**

The function of the translocation domain and the ER retention domain can be tested as a function of the chimera's ability to gain access to the cytosol. Because access first requires binding to the cell, these assays also are useful to determine whether the cell recognition domain is functioning.

a. **Presence in the cytosol**

In one method, access to the cytosol is determined by detecting the physical presence of the chimera in the cytosol. For example, the chimera can be labelled and the chimera exposed to the cell. Then, the cytosolic fraction is isolated and the amount of label in the fraction determined. Detecting label in the fraction indicates that the chimera has gained access to the cytosol.

b. **ADP Ribosylation activity**

In another method, the ability of the translocation domain and ER retention domain to effect translocation to the cytosol can be tested with a construct containing a

domain III having ADP ribosylation activity. Briefly, cells are seeded in tissue culture plates and exposed to the chimeric protein or to an engineered PE exotoxin containing the modified translocation domain or ER retention sequence in place of the native domains. ADP ribosylation activity is determined as a function of inhibition of protein synthesis by, e.g., monitoring the incorporation of ^3H -leucine.

3. Immunogenicity

The function of the non-native epitope can be determined by determining humoral or cell-mediated immunogenicity. Immunogenicity can be tested by several methods. Humoral immune response can be tested by inoculating an animal and detecting the production of antibodies against the foreign immunogen. Cell-mediated cytotoxic immune responses can be tested by immunizing an animal with the immunogen, isolating cytotoxic T cells, and detecting their ability to kill cells whose MHC Class I molecules bear amino acid sequences from the non-native epitope. Because generating a cytotoxic T cell response requires both binding of the chimera to the cell and translocation to the cytosol, this test also tests the activity of the cell recognition domain, the translocation domain and the ER retention domain.

III. RECOMBINANT POLYNUCLEOTIDES ENCODING PE-LIKE CHIMERIC IMMUNOGENS

A. Recombinant Polynucleotides

1. Sources

This invention provides recombinant polynucleotides comprising a nucleotide sequence encoding the PE-like chimeric immunogens of this invention. These polynucleotides are useful for making the PE-like chimeric immunogens. In another aspect, this invention provides a PE-like protein cloning platform comprising a recombinant polynucleotide sequence encoding a cell recognition domain, a translocation domain, an ER retention domain and, between the translocation domain and the ER retention domain, a cloning site for a polynucleotide sequence encoding a non-native epitope domain.

The recombinant polynucleotides of this invention are based on polynucleotides encoding *Pseudomonas* exotoxin A, or portions of it. A nucleotide sequence encoding PE is presented above. The practitioner can use this sequence to

prepare PCR primers for isolating a full-length sequence. The sequence of PE can be modified to engineer a polynucleotide encoding the PE-like chimeric immunogen or platform.

5 A polynucleotide encoding PE or any other polynucleotide used in the chimeric proteins of the invention can be cloned or amplified by *in vitro* methods, such as the polymerase chain reaction (PCR), the ligase chain reaction (LCR), the transcription-based amplification system (TAS), the self-sustained sequence replication system (3SR) and the Q β replicase amplification system (QB). For example, a polynucleotide encoding the protein can be isolated by polymerase chain reaction of
10 cDNA using primers based on the DNA sequence of PE or a cell recognition molecule.

A wide variety of cloning and *in vitro* amplification methodologies are well-known to persons skilled in the art. PCR methods are described in, for example, U.S. Pat. No. 4,683,195; Mullis *et al.* (1987) *Cold Spring Harbor Symp. Quant. Biol.* 51:263; and Erlich, ed., *PCR Technology*, (Stockton Press, NY, 1989). Polynucleotides
15 also can be isolated by screening genomic or cDNA libraries with probes selected from the sequences of the desired polynucleotide under stringent hybridization conditions.

2. Mutagenized versions

Mutant versions of the proteins can be made by site-specific mutagenesis of other polynucleotides encoding the proteins, or by random mutagenesis caused by increasing the error rate of PCR of the original polynucleotide with 0.1 mM MnCl₂ and unbalanced nucleotide concentrations.

Eliminating nucleotides encoding amino acids 1-252 yields a construct referred to as "PE40." Eliminating nucleotides encoding amino acids 1-279 yields a construct referred to as "PE37." (See Pastan *et al.*, U.S. patent 5,602,095.) The practitioner can ligate sequences encoding cell recognition domains to the 5' end of these platforms to engineer PE-like chimeric proteins that are directed to particular cell surface receptors. These constructs optionally can encode an amino-terminal methionine. A cell recognition domain can be inserted into such constructs in the nucleotide sequence encoding the ER retention domain.

3. Chimeric protein cloning platforms

A cloning site for the non-native epitope domain can be introduced between the nucleotides encoding the cysteine residues of domain Ib. For example, as described in the Examples, a nucleotide sequence encoding a portion of the Ib domain between the cysteine-encoding residues can be removed and replaced with a nucleotide sequence encoding an amino acid sequence and that includes a PstI cloning site. A polynucleotide encoding the non-native epitope and flanked by PstI sequences can be inserted into the vector.

The construct also can be engineered to encode a secretory sequence at the amino terminus of the protein. Such constructs are useful for producing the immunogens in mammalian cells. *In vitro*, such constructs simplify isolation of the immunogen. *In vivo*, the constructs are useful as polynucleotide vaccines; cells that incorporate the construct will express the protein and secrete it where it can interact with the immune system.

B. Expression Vectors

This invention also provides expression vectors for expressing PE-like chimeric immunogens. Expression vectors are recombinant polynucleotide molecules comprising expression control sequences operatively linked to a nucleotide sequence encoding a polypeptide. Expression vectors can be adapted for function in prokaryotes

or eukaryotes by inclusion of appropriate promoters, replication sequences, markers, etc. for transcription and translation of mRNA. The construction of expression vectors and the expression of genes in transfected cells involves the use of molecular cloning techniques also well known in the art. Sambrook *et al.*, *Molecular Cloning -- A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, (1989) and *Current Protocols in Molecular Biology*, F.M. Ausubel *et al.*, eds., (Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc.) Useful promoters for such purposes include a metallothionein promoter, a constitutive adenovirus major late promoter, a dexamethasone-inducible MMTV promoter, a SV40 promoter, a MRP polIII promoter, a constitutive MPSV promoter, a tetracycline-inducible CMV promoter (such as the human immediate-early CMV promoter), and a constitutive CMV promoter. A plasmid useful for gene therapy can comprise other functional elements, such as selectable markers, identification regions, and other genes.

Expression vectors useful in this invention depend on their intended use. Such expression vectors must, of course, contain expression and replication signals compatible with the host cell. Expression vectors useful for expressing PE-like chimeric immunogens include viral vectors such as retroviruses, adenoviruses and adeno-associated viruses, plasmid vectors, cosmids, and the like. Viral and plasmid vectors are preferred for transfecting mammalian cells. The expression vector pcDNA1 (Invitrogen, San Diego, CA), in which the expression control sequence comprises the CMV promoter, provides good rates of transfection and expression. Adeno-associated viral vectors are useful in the gene therapy methods of this invention.

A variety of means are available for delivering polynucleotides to cells including, for example, direct uptake of the molecule by a cell from solution, facilitated uptake through lipofection (e.g., liposomes or immunoliposomes), particle-mediated transfection, and intracellular expression from an expression cassette having an expression control sequence operably linked to a nucleotide sequence that encodes the inhibitory polynucleotide. See also Inouye *et al.*, U.S. Patent 5,272,065; *Methods in Enzymology*, vol. 185, Academic Press, Inc., San Diego, CA (D.V. Goeddel, ed.) (1990) or M. Krieger, *Gene Transfer and Expression -- A Laboratory Manual*, Stockton Press, New York, NY, (1990). Recombinant DNA expression plasmids can also be used to prepare the polynucleotides of the invention for delivery by means other than by gene

therapy, although it may be more economical to make short oligonucleotides by *in vitro* chemical synthesis.

The construct can also contain a tag to simplify isolation of the protein. For example, a polyhistidine tag of, e.g., six histidine residues, can be incorporated at the amino terminal end of the protein. The polyhistidine tag allows convenient isolation of the protein in a single step by nickel-chelate chromatography.

C. Recombinant Cells

The invention also provides recombinant cells comprising an expression vector for expression of the nucleotide sequences encoding a PE chimeric immunogen of this invention. Host cells can be selected for high levels of expression in order to purify the protein. The cells can be prokaryotic cells, such as *E. coli*, or eukaryotic cells. Useful eukaryotic cells include yeast and mammalian cells. The cell can be, e.g., a recombinant cell in culture or a cell *in vivo*.

E. coli has been successfully used to produce PE-like chimeric immunogens. The protein can fold and disulfide bonds can form in this cell.

IV. PSEUDOMONAS EXOTOXIN A-LIKE CHIMERIC IMMUNOGEN VACCINES

PE-like chimeric immunogens are useful in vaccines for eliciting a protective immune response against agents bearing the non-native epitope. A vaccine can include one or a plurality (i.e. a multivalent vaccine) of different PE-like chimeric immunogens. For example, a vaccine can include PE-like chimeric immunogens whose non-native epitopes come from several circulating strains of a pathogen. As the pathogen evolves, new PE-like chimeric immunogens can be constructed that include the altered epitopes, for example, from breakthrough viruses. In one embodiment, the vaccine comprises epitopes from a T-cell-tropic virus and from a macrophage-tropic virus. For example, a vaccine against HIV infection can include immunogens whose non-native epitopes derive from the V3 loop of MN and Thai-E strains of HIV. Also, the epitopes can derive from any peptide from HIV that is involved in membrane fusion, e.g., gp120 or gp41. Alternatively, because they are subunit vaccines, the vaccine can include PE-like chimeric immunogens whose non-native epitopes are selected from various epitopes of the same pathogen.

The vaccine can come lyophilized or already reconstituted in sterile solution for use. An immunizing dose is between about 1 μg and about 1000 μg , more usually between about 10 μg and about 50 μg of the recombinant protein. For determination of immunizing doses see, for example, *Manual of Clinical Immunology*, H.R. Rose and H. Friedman, American Society for Microbiology, Washington, D.C. (1980). A unit dose is about 0.05 ml to about 1 ml, more usually about 0.5 ml. A dose is preferably delivered subcutaneously or intramuscularly. An injection can be followed by several more injections spaced about 4 to about 8 weeks apart. Booster doses can follow in about 1 to about 10 years. The vaccine can be prepared in dosage forms containing between 1 and 50 doses (e.g., 0.5 ml to 25 ml), more usually between 1 and 10 doses (e.g., 0.5 ml to 5 ml). The vaccine also can include an adjuvant, that potentiates an immune response when used in conjunction with an antigen. Useful adjuvants include alum, aluminum hydroxide or aluminum phosphate.

V. METHODS OF ELICITING AN IMMUNE RESPONSE

PE-like chimeric immunogens are useful in eliciting an immune response against antigens bearing the non-native epitope. Eliciting a humoral immune response is useful in the production of antibodies that specifically recognize the non-native epitope and in immunization against cells, viruses or other agents that bear the non-native epitope. PE-like chimeric immunogens are also useful in eliciting MHC Class I-dependent or MHC Class II-dependent cell-mediated immune responses. They are also useful in eliciting a secretory immune response.

A. Prophylactic and Therapeutic Treatments

PE-like chimeric immunogens can include non-native epitopes from pathogenic organisms or from pathological cells from a subject, such as cancer cells. Accordingly, this invention provides prophylactic and therapeutic treatments for diseases involving the pathological activity of agents, either pathogens or aberrant cells, that bear the non-native epitope. The methods involve immunizing a subject with PE-like chimeric immunogens bearing the non-native epitope. The resulting immune responses mount an attack against the pathogens, themselves, or against cells that express the non-native epitope. For example, if the pathology results from bacterial or parasitic protozoan infection, the immune system mounts a response against the pathogens, themselves. If

the pathogen is a virus, infected cells will express the non-native epitope on their surface and become the target of a cytotoxic response. Aberrant cells, such as cancer cells, often express un-normal epitopes, and also can be subject to a cytotoxic immune response.

B. Humoral Immune Response

PE-like chimeric immunogens are useful in eliciting the production of antibodies against the non-native epitope by a subject. PE-like chimeric immunogens are attractive immunogens for making antibodies against non-native epitopes that naturally occur within a cysteine-cysteine loop: Because they contain the non-native epitope within a cysteine-cysteine loop, they present the epitope to the immune system in near-native conformation. The resulting antibodies generally recognize the native antigen better than those raised against linearized versions of the non-native epitope.

Methods for producing polyclonal antibodies are known to those of skill in the art. In brief, an immunogen, preferably a purified polypeptide, a polypeptide coupled to an appropriate carrier (e.g., GST, keyhole limpet hemanocyanin, etc.), or a polypeptide incorporated into an immunization vector, such as a recombinant vaccinia virus (see, U.S. Patent No. 4,722,848) is mixed with an adjuvant. Animals are immunized with the mixture. An animal's immune response to the immunogenic preparation is monitored by taking test bleeds and determining the titer of reactivity to the polypeptide of interest. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the polypeptide is performed where desired. See, e.g., Coligan (1991) *Current Protocols in Immunology* Wiley/Greene, NY; and Harlow and Lane (1989) *Antibodies: A Laboratory Manual* Cold Spring Harbor Press, NY.

In various embodiments, the antibodies ultimately produced can be monoclonal antibodies, humanized antibodies, chimeric antibodies or antibody fragments.

Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies are screened for binding to polypeptides comprising the epitope, or screened for agonistic or antagonistic activity, e.g., activity mediated through the agent comprising the non-native epitope. In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates,

humans, etc. Description of techniques for preparing such monoclonal antibodies are found in, e.g., Stites *et al.* (eds.) *Basic and Clinical Immunology* (4th ed.) Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane, *Supra*; Goding (1986) *Monoclonal Antibodies: Principles and Practice* (2d ed.) Academic Press, New York, NY; and Kohler and Milstein (1975) *Nature* 256: 495-497.

In another embodiment, the antibodies are humanized immunoglobulins. Humanized antibodies are made by linking the CDR regions of non-human antibodies to human constant regions by recombinant DNA techniques. See Queen *et al.*, United States patent 5,585,089.

In another embodiment of the invention, fragments of antibodies against the non-native epitope are provided. Typically, these fragments exhibit specific binding to the non-native epitope similar to that of a complete immunoglobulin. Antibody fragments include separate heavy chains, light chains, Fab, Fab' F(ab')₂ and Fv. Fragments are produced by recombinant DNA techniques, or by enzymic or chemical separation of intact immunoglobulins.

Other suitable techniques involve selection of libraries of recombinant antibodies in phage or similar vectors. See, Huse *et al.* (1989) *Science* 246: 1275-1281; and Ward *et al.* (1989) *Nature* 341: 544-546.

An approach for isolating DNA sequences which encode a human monoclonal antibody or a binding fragment thereof is by screening a DNA library from human B cells according to the general protocol outlined by Huse *et al.*, *Science* 246:1275-1281 (1989) and then cloning and amplifying the sequences which encode the antibody (or binding fragment) of the desired specificity. The protocol described by Huse is rendered more efficient in combination with phage display technology. See, e.g., Dower *et al.*, WO 91/17271 and McCafferty *et al.*, WO 92/01047. Phage display technology can also be used to mutagenize CDR regions of antibodies previously shown to have affinity for the polypeptides of this invention or their ligands. Antibodies having improved binding affinity are selected.

The antibodies of this invention are useful for affinity chromatography in isolating agents bearing the non-native epitope. Columns are prepared, e.g., with the antibodies linked to a solid support, e.g., particles, such as agarose, Sephadex, or the like, where a cell lysate is passed through the column, washed, and treated with increasing concentrations of a mild denaturant, whereby purified agents are released.

Antibodies were produced against gp120 using a PE-like chimeric immunogen having the gp120 V3 loop as the non-native epitope. The monoclonal antibodies selectively captured the soluble MN and Th-E chimeric proteins, confirming that the V3 loops were exposed and accessible to antibody probes. Also, sera from immunized rabbits neutralized HIV-1 infectivity in an *in vitro* assay.

C. MHC Class II-dependent Cell-mediated Immune Response

In another aspect, this invention provides methods for eliciting an MHC Class II-dependent immune response against cells expressing the non-native epitope.

MHC Class II molecules bind peptides having particular amino acid motifs well known in the art. The MHC Class II-dependent response involves the uptake of an antigen by antigen-presenting cells (APC's), its processing, and presentation on the cell surface as part of an MHC Class II/antigenic peptide complex. Alternatively, MHC Class II molecules on the cell surface can bind peptides having the proper motif.

Antigen presenting cells interact with CD4-positive T-helper cells, thereby activating the T-helper cells. Activated T-helper cells stimulate B-lymphocytes to produce antibodies against the antigen. Antibodies mark cells bearing the antigen on their surface. The marked cells are subject to antibody-dependent cell-mediated cytotoxicity, in which NK cells or macrophages, which bear Fc receptors, attack the marked cells.

Methods for eliciting an MHC Class II-dependent immune response involve administering to a subject a vaccine including an immunogenic amount of a chimeric *Pseudomonas* exotoxin that includes an amino acid motif recognized by MHC Class II molecules of the subject. Alternatively, antigen presenting cells can be cultured with such peptides to allow binding, and the cells can be administered to the subject. Preferably, the cells are syngeneic with the subject.

D. MHC Class I-dependent Cell-mediated Immune Response

In another aspect, this invention provides methods for eliciting an MHC Class I-dependent cell-mediated immune response against cells expressing the non-native epitope in a subject. MHC Class I molecules bind peptides having particular amino acid motifs well known in the art. Proteins expressed in a cell are digested into peptides and presented on the cell surface in association with MHC Class I molecules. There, they

are recognized by CD8-positive lymphocytes, generating a cytotoxic T-lymphocyte response against cells expressing the epitopes in association with MHC Class I molecules. Because CD4-positive T lymphocytes infected with HIV express gp120 and, thus, the V3 domain, the generation of cytotoxic T-lymphocytes that attack such cells is useful in the prophylactic or therapeutic treatment of HIV infections.

HLA-A1 binding motif includes a first conserved residue of T, S or M, a second conserved residue of D or E, and a third conserved residue of Y. Other second conserved residues are A, S or T. The first and second conserved residues are adjacent and are preferably separated from the third conserved residue by 6 to 7 residues. A second motif consists of a first conserved residue of E or D and a second conserved residue of Y where the first and second conserved residues are separated by 5 to 6 residues. The HLA-A3.2 binding motif includes a first conserved residue of L, M, I, V, S, A, T and F at position 2 and a second conserved residue of K, R or Y at the C-terminal end. Other first conserved residues are C, G or D and alternatively E. Other second conserved residues are H or F. The first and second conserved residues are preferably separated by 6 to 7 residues. The HLA-A11 binding motif includes a first conserved residue of T or V at position 2 and a C-terminal conserved residue of K. The first and second conserved residues are preferably separated by 6 or 7 residues. The HLA-A24.1 binding motif includes a first conserved residue of Y, F or W at position 2 and a C terminal conserved residue of F, I, W, M or L. The first and second conserved residues are preferably separated by 6 to 7 residues.

Another method involves transfecting cells *ex vivo* with such expression vectors, and administering the cells to the subject. The cells preferably are syngeneic to the subject.

Methods for eliciting an immune response against a virus in a subject are useful in prophylactic methods for preventing infection with the virus when the vaccine is administered to a subject who is not already infected.

E. IgA-mediated Secretory Immune Response

Mucosal membranes are primary entryways for many infectious pathogens. Such pathogens include, for example, HIV, herpes, vaccinia, cytomegalovirus, yersinia and vibrio. Mucosal membranes include the mouth, nose, throat, lung, vagina, rectum and colon. As a defense against entry, the body secretes secretory IgA on the surfaces

of mucosal epithelial membranes against pathogens. Furthermore, antigens presented at one mucosal surface can trigger responses at other mucosal surfaces due to trafficking of antibody-secreting cells between these mucosae. The structure of secretory IgA has been suggested to be crucial for its sustained residence and effective function at the luminal surface of a mucosa. As used herein, "secretory IgA" or "sIgA" refers to a polymeric molecule comprising two IgA immunoglobulins joined by a J chain and further bound to a secretory component. While mucosal administration of antigens can generate an IgG response, parenteral administration of immunogens rarely produce strong sIgA responses. Generating a secretory immune response for defense against HIV is a recognized need. (Bukawa, H., *et al.* 1995, *Nat Med* 1, 681-5; Mestecky, J., *et. al.*, 1994, *Aids Res Hum Retroviruses* 10, S11-20.)

Pseudomonas exotoxin binds to receptors on mucosal membranes. Therefore, PE-like chimeric immunogens are an attractive vector for bringing non-native epitopes to a mucosal surface. There, the immunogens elicit an IgA-mediated immune response against the immunogen. Accordingly, this invention provides PE-like chimeric immunogens comprising a non-native epitope from a pathogen that gains entry through mucosal membranes. The cell recognition domain can be targeted to any mucosal surface receptor. These PE-like chimeric immunogens are useful for eliciting an IgA-mediated secretory immune response against immunogens that gain entry to the body through mucosal surfaces. PE-like chimeric immunogens used for this purpose should have ligands that bind to receptors on mucosal membranes as their cell recognition domains. For example, epidermal growth factor binds to the epidermal growth factor receptor on mucosal surfaces.

The immunogens can be applied to the mucosal surface by any of the typical means, including pharmaceutical compositions in the form of liquids or solids, e.g., sprays, ointments, suppositories or erodible polymers impregnated with the immunogen. Administration can involve applying the immunogen to a plurality of different mucosal surfaces in a series of immunizations, e.g., as booster immunizations. A booster inoculation also can be administered parenterally, e.g., subcutaneously. The immunogen can be administered in doses of about 1 μ g to 1000 μ g, e.g., about 10 μ g to 100 μ g.

Subcutaneous inoculation with vaccines comprising an epitope from the principal neutralizing domain of gp120 of HIV is not known to generate secretory IgA.

Accordingly, mucosal presentation of the chimeric immunogens of this invention is useful for producing these hitherto unknown antibodies. This invention also provides secretory IgA that specifically recognize epitopes of other pathogens that enter the body through a mucous membrane.

5 The IgA response is strongest on mucosal surfaces exposed to the immunogen. Therefore, in one embodiment, the immunogen is applied to a mucosal surface that is likely to be a site of exposure to the particular pathogen. Accordingly, chimeric immunogens against sexually transmitted diseases can be administered to vaginal, anal or oral mucosal surfaces.

10 Mucosal administration of the chimeric immunogens of this invention result in strong memory responses, both for IgA and IgG. Therefore, in vaccination with them, it is useful to provide booster doses either mucosally or parenterally. The memory response can be elicited by administering a booster dose more than a year after the initial dose. For example, a booster dose can be administered about 12, about 16,
15 about 20 or about 24 months after the initial dose.

VI. POLYNUCLEOTIDE VACCINES AND METHODS OF GENE THERAPY

Vaccines comprising polynucleotides encoding a protein immunogen, often called "DNA vaccines," offer certain advantages over polypeptide vaccines. DNA
20 vaccines do not run the risk of contamination with unwanted protein immunogens. Upon administration to a subject, the polynucleotide is taken up by a cell. RNA is reverse transcribed into DNA. DNA is integrated into the genome in some percentage of transfected cells. Where the DNA integrates so as to be operatively linked with expression control sequences, or if such sequences are provided with the recombinant
25 polynucleotide, the cell expresses the encoded polypeptide. Upon secretion from the cell, the polypeptide acts as an immunogen. Naked DNA is preferentially taken up by liver and by muscle cells. Accordingly, the polypeptide can be injected into muscle tissue, or provided by, e.g., biolistic injection. Generally, doses of naked polynucleotide will be from about 1 μ g to 100 μ g for a typical 70 kg patient.

30 The polynucleotide vaccines of this invention can include polynucleotides encoding PE-like chimeric immunogens that are used in polypeptide vaccines. This includes multiple immunogens including several variants of an epitope.

The following examples are offered by way of illustration, not by way of limitation.

EXAMPLES

I. CONSTRUCTION OF PE-LIKE CHIMERIC IMMUNOGENS

To generate chimeric proteins, the subdomain Ib of ntPE was replaced with V3 loop sequences from either an MN (subtype B) or Thai-E subtype strain of HIV-1. The MN sequence is from a T-cell-tropic strain while the Thai-E sequence comes from a macrophage-tropic strain.

Wild-type (WT) PE is composed of 613 amino acids and has a molecular mass of 67,122 Da. Deletion of a glutamic acid 553 ($\Delta E553$) results in a non-toxic version of PE (Lukac, M., *et al.*, 1988, *Infect and Immun* 56:3095-3098), referred to as ntPE.

Plasmids were constructed by inserting oligonucleotide duplexes encoding V3 loop sequences into a new PE-based vector that was designed with a novel PstI site. In an effort to produce a V3 loop of similar topology to that found in gp120, the 14 or 26 amino acid inserts were flanked by cysteine residues (Fig. 1C - bold type). Construction of the novel vector resulted in several changes in the amino acid sequence of ntPE near the insertion point of the V3 loop (Fig. 1C - italics). The non-toxic chimeras, ntPE-V3MN14, ntPE-V3MN26 and ntPE-V3Th-E26, contained V3 loops of 14 or 26 amino acids from the MN strain or 26 amino acids from the Thai-E strain, respectively (nt = "non-toxic"). Insertion of an irrelevant 16 amino acid sequence resulted in the construction of a control chimera referred to as ntPE-fp126. Removal of the Ib loop (6 amino acids) and modification of flanking amino acids adjacent to the V3 loop insert resulted in a small increase in molecular mass compared to wild-type PE (Fig. 1C).

More specifically, plasmid pMOA1A2VK352 (Ogata, M., *et al.*, 1992, *J Biol Chem*, 267, 25396-401), encoding PE, was digested with SfiI and ApaI (residues 1143 and 1275, respectively) and then re-ligated with a duplex containing a novel PstI site. The coding strand of the duplex had the following sequence: 5'-tggccctgac cctggccgcc gccgagagcg agcgcttcgt ccggcagggc accggcaacg acgaggccgg cgcggcaaac ctgcagggcc -3' (SEQ ID NO:5). The resulting plasmid encoded a slightly smaller version of PE and lacked much of domain Ib. The PstI site was then used to introduce

duplexes encoding V3 loop sequences flanked by cysteine residues. To make non-toxic proteins, vectors were modified by the subcloning in an enzymatically inactive domain III from pVC45ΔE553. An additional subcloning, from pJH4 (Hwang, J., *et. al.*, 1987, *Cell*, 48, 129-136), was needed to produce a vector that lacked a signal sequence.

Insertion of duplexes and subcloning modifications were initially verified by restriction analysis while final constructs were confirmed by dideoxy double strand sequencing.

II. CHARACTERIZATION OF CHIMERAS

A. Expression

All ntPE-V3 loop chimeric proteins were expressed in *E. coli* SA2821/BL21(ΔDE3) using a T7 promoter/T7 polymerase system (Studier, F.W., *et. al.*, 1990, *Methods Enzymol* 185, 60-89). SA2821/BL21(ΔDE3) cells were transformed with the appropriate plasmid and grown to an absorbance of 1.0 (600 nm) in medium containing ampicillin. To induce high level protein expression, isopropyl-β-D-thiogalactoside (1 mM) was added to the culture and incubated for an additional 90 min. *E. coli* cell pellets, were resuspended in 50 mM Tris/20 mM EDTA, pH 8.0 (TE buffer) and dispersed using a Tissue Miser. Cell lysis was accomplished with lysozyme (200 μg/ml final concentration; Sigma) and membrane associated proteins were solubilized by the addition of 2.5% Triton X-100 and 0.5 M NaCl.

PE-V3 loop chimeras were present in inclusion bodies, which were recovered by centrifugation. After washing with TE containing 0.5% Triton X-100 and then with TE alone, inclusion bodies were solubilized by the addition of 6 M guanidine and 65 mM dithioerythritol. Refolding was allowed to proceed at a final protein concentration of 100 μg/ml for a minimum of 24 h at 8°C in 0.1 M Tris (pH 8.0) containing 0.5 M L-arginine (Sigma), 2 mM EDTA and 0.9 mM glutathione. The protease inhibitor AEBSF (Boehringer Mannheim) was added to a final concentration of 0.5 mM. Proteins were dialyzed against 20 mM Tris, 2 mM EDTA and 100 mM urea, pH 7.4. Following dialysis, proteins were applied to a Q sepharose column (Pharmacia Biotech; Piscataway, NJ). After washing with 20 mM Tris (pH 8-0) containing 0.1 M NaCl, chimeric proteins were eluted with 0.3 M NaCl in the same buffer and concentrated using Centriprep-30 ultrafiltration devices (Amicon, Inc.; Beverly, MA). An HPLC gel filtration column (G3000SW from Toso Haas; Montgomeryville, PA) was

used to isolate final products. A typical yield of properly folded protein per 4L bacterial culture was 50-100 mg with a purity greater than 95%.

B. Biochemical Characterization

Chimeric proteins were separated by SDS-PAGE using 8-16% gradient polyacrylamide gels (Novex; San Diego, CA), and visualized by staining with Coomassie Blue. For Western blot analysis, proteins were transferred onto Immobilon-P membranes (Millipore Corp., Bedford, MA) and exposed to either an anti-PE mouse monoclonal antibody (M40-1 (Ogata, M., *et. al.*, 1991, *Infect and Immun* 59, 407-414) or an anti-gp120 mouse monoclonal antibody (1F12 for MN sequences or 1B2 for Thai-E sequences; Genentech, Inc.; South San Francisco, CA). The primary antibody was detected by a secondary anti-mouse antibody conjugated to horseradish peroxidase. Reactive products were visualized by the addition of diaminobenzadine and hydrogen peroxide. Immunocapture experiments were performed for 30 min at 23°C using the 1F12 anti-gp120 monoclonal antibody. Antibody-chimeric protein complexes were recovered with protein G sepharose beads (Pharmacia Biotech; Piscataway, NJ) and separated using SDS-PAGE (as above). Recombinant forms of gp120 derived from HIV-1-MN (120/MN; Genentech, Inc.) and the Thai subtype E isolate (gp120/Th-E - Chiang Mai; Advanced Biotechnologies, Columbia MD) were used as standards.

SDS-PAGE analysis of purified ntPE-V3 loop chimeras (Fig. 2A) was consistent with calculated masses (Fig. 1C). Western blots, using monoclonal antibodies raised against gp120/MN (1F12) or gp120/Th-E (1B2), showed strain-specific reactivity with the MN and Thai-E V3 loop chimeras (Fig. 2B).

Free sulfhydryl analysis of purified ntPE-V3 loop chimeras failed to demonstrate any unpaired cysteines, suggesting that the purified ntPE-V3 loop chimeras had refolded and oxidized to form a disulfide bond at the base of the V3 loop (Fig. 1A). The formation of this disulfide bond was expected to result in the exposure of the V3 loop at the surface of the chimeras.

To determine sulfhydryl content, chimeric proteins (15 nmols) in PBS (pH 7.4) containing 1 mM EDTA, were reacted with 1 mM thionitrobenzoate (DTNB) (Pierce Chem Co, Rockford, IL) for 15 min at 23°C. The release of thionitrobenzoate was monitored at 412 nm. DTNB reactivity was confirmed by the use of cysteine.

This was tested directly by immuno-capture studies (Fig. 2C). The 1F12 and 1B2 monoclonal antibodies selectively captured the soluble MN and Th-E chimeric proteins confirming that the V3 loops were exposed and accessible to antibody probes. Despite the fact that the 1F12 antibody reacted strongly with ntPE-V3MN14 in Western blots (Fig 2B), it captured only a small amount of soluble protein (Fig. 2C, Lane 3), suggesting that the reactive epitope was not completely exposed when only 14 amino acids were inserted.

C. Circular Dichroism

To evaluate the impact of amino acid inserts on the secondary structure of the chimeras, near- and far-UV CD spectral analysis was performed on purified ntPE-V3MN14 and ntPE-V3MN26 proteins and compared these to wild-type PE (wtPE) spectra (Figs. 3A and 3B). Circular dichroism (CD) spectra were collected on an Aviv 60DS spectropolarimeter. Near UV CD spectra (400 nm to 250 nm) were obtained in 0.2 nm increments with a 0.5 nm bandwidth and a 5 second time constant (150 readings/second averaged) for samples in a 1 cm pathlength cell. Far UV spectra (250 nm to 190 nm) were collected in 0.2 nm increments with a 0.5 nm bandwidth and a 3 second time constant in a 0.05 cm pathlength cell. Each spectrum was digitally smoothed using the Savitsky-Golay algorithm (Gorry, P.A. 1990, *Analytical Chem* 62, 570-573), corrected for concentration, and normalized to units of mean residue weight ellipticity (Θ MRW) using the following relationship:

$$\theta_{MRW} = \frac{\theta_{obs} (MW_{monomer} / n_{monomer})}{10 (d) (c)}$$

where Θ_{obs} is the observed ellipticity, $MW_{monomer}$ is the molecular weight of the monomer, $n_{monomer}$ is the number of amino acids in the monomer, d is the pathlength of the cell (cm), and c is the concentration of the sample in the cell (mg/ml).

Secondary structure calculations (Fig. 3C) suggested that there were no significant differences between these proteins and wtPE. ntPE-V3MN14 demonstrated more negative ellipticity than ntPE-V3MN26 and wtPE, suggesting more strain may occur on the disulfide bond at the base of the loop insert for this chimera. Both ntPE-V3MN14 and ntPE-V3MN26 showed an apparent red-shift at 290 nm, possibly due to the additional tyrosine residues in the chimeras. Alternately, this red-shift could result

from a slight environmental perturbation of a tryptophan residue. Altogether, these results suggest that the V3 loop inserts did not produce large alterations in the secondary structure relative to wild-type toxin and that the changes in tertiary structure were consistent with the presence of the 14 and 26 amino acid inserts.

III. TRANSLOCATION TO THE CYTOSOL

After binding to the LRP receptor, ntPE-V3 loop chimeras should be endocytosed, cleaved by furin and the C-terminal portion containing domains II, the V3 loop and III should be translocated to the cytosol in a similar fashion to wtPE (Ogata, M., *et. al.*, 1990, *Biol Chem* 265, 20678-85). This was tested directly by producing enzymatically active versions of PE-V3MN14 and 26 (containing glutamic acid 553 and having the ability to ADP-ribosylate elongation factor 2) and comparing their activity with wtPE in cytotoxicity assays.

Human A431 (epidermoid carcinoma) cells were seeded in 24-well tissue culture plates at 1×10^5 cells/well in RPMI 1640 media supplemented with 5% fetal bovine serum. After 24 h, cells were treated for 18 h at 37°C with 4-fold dilutions of either wtPE or toxic forms (with a glutamic acid residue at position 553 and capable of ADP-ribosylating elongation factor 2) of the chimeric proteins. Inhibition of protein synthesis was assessed by monitoring the incorporation of ^3H -leucine.

When assayed for its ability to inhibit protein synthesis, PE-V3MN26 exhibited similar toxicity to wtPE in human A431 cells (Fig. 4). PE-V3MN14 was also fully toxic. These results confirmed that the size and location of the V3 loop inserts did not impede toxin delivery to the cytosol. Further, these data suggest that the isolation, refolding and purification protocol used to prepare these chimeras resulted in the production of a correctly folded and functional protein.

IV. IMMUNOGENICITY

To investigate their usefulness as immunogens, rabbits were injected subcutaneously with 200 μg of either the MN or Thai-E chimeras. Rabbits were immunized subcutaneously at four sites with 200 μg (total) of ntPE-V3MN26. The first injection was administered with complete Freund's adjuvant. All subsequent injections (at 2, 4 and 12 weeks) were given with incomplete Freund's adjuvant. Venous bleeds

were obtained weekly after the third injection and screened by immunoblotting against gp120.

In Western blots, serum samples from rabbits immunized with the ntPE-V3MN proteins exhibited a strong reactivity for immobilized recombinant gp120/MN (Fig. 5A). Reactive titers increased with time: at 6 weeks reactivity was noted at 1:200 dilution, at 12 weeks at 1:5,000 dilution and at later times reactivity could be detected at 1:25,000. These anti-V3 loop/MN sera were not reactive with gp120/Thai-E (Fig. 5A). Sera from rabbits injected with non-toxic PE (i.e. ntPE with no insert) exhibited no reactivity for gp120. Rabbits injected with the ntPE-V3Th-E produced reactive sera for gp120/Thai-E but not for gp120/MN (Fig. 5A).

Sera from rabbits immunized with ntPE-V3MN26 were characterized further. Reactivity for immobilized gp120/MN was absorbed when these sera were pre-mixed with soluble recombinant gp120/MN (Fig. 5B). This blocking activity, which was dose-dependent and maximal at 50 µg/ml, indicated that rabbits responded primarily to V3 loop sequences that are exposed on the surface of gp120.

Sera from immunized rabbits were also found to neutralize HIV-1 infectivity in an *in vitro* assay (Fig. 6). This assay utilized MT4 cells as an indicator of HIV-1-mediated cell death (Miyoshi, I., *et al.*, 1981, *Nature* 294, 770-1). Duplicate serial dilutions of antiserum was incubated with HIV-1/MN grown in FDA/H9 cells (Popovic, M., *et al.*, 1984, *Science* 224, 497-500) and the mixture added to cells for 7 days. Viral-mediated cell death was assessed using a MTT dye assay (Robertson, G.A., *et al.*, 1988, *J Virol Methods* 20, 195-202) and spectrophotometric analysis at 570 nm. The serum 50% inhibitory concentration was calculated and reported as the neutralization titer.

Pre-immune sera did not show any protection of a human T-cell line, MT4, from killing by HIV-1 MN. Although sera at 5 weeks following immunization also showed no protection, week 8 and week 27 sera were protective against viral challenge with 50% neutralization occurring at approximately a 1:400 dilution. Based upon the immunization schedule used, week 5 sera reflected the response in animals immunized and boosted once, while week 8 sera was from animals boosted twice and week 27 sera came from animals boosted three times. MT4 cell survival values obtained for sera dilutions of less than 1:100 for the week 8 and week 27 bleeds were greater than the unchallenged cell control used for normalization. This was likely due to stimulation

by growth factors present in the rabbit sera. The data suggest that the immune response following subcutaneous injections of ntPE-V3 loop chimeras can result in the production of neutralizing antibodies.

5 V. NEUTRALIZATION OF INFECTIVITY

Antibodies elicited by the chimeric immunogen were shown to have the ability to neutralize infectivity of HIV-1 in viral growth assays where suppression of p24 production was used as an indicator of HIV neutralization. Clinical isolates corresponding to subtype B, RVL05, and subtype E, Th92009, were incubated with
10 dilutions of rabbit sera and cultured in PBMCs for a total of 5 days.

One assay utilized MT4 cells as an indicator of HIV-1-mediated cell death. I. Miyoshi et al. (1981) *Nature* 294:770-771. Duplicate serial dilutions of antiserum were incubated with HIV-1/MN and grown in FDA/H9 cells and the mixture added to MT4 cells for 7 days. M. Popovic et al. (1984) *Science* 224:497-500. Viral-mediated
15 cell death was assessed using a 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide dye assay and spectrophotometric analysis at 570 nm. G.A. Robertson et al. (1988) *J. Virol. Methods* 20:195-202. The serum 50% inhibitory concentration was calculated and reported as the neutralization titer.

A second assay used p24 production of as an indicator of viral growth. T. Wrin et al. (1995) *J. Virol.* 69:39-48. Primary virus was first titrated to determine the amount that reproducibly yielded significant but submaximal amounts of p24. Virus
20 preparations were incubated for 1 h at 37° C with various dilutions of rabbit sera, either immune or pre-bleed, and this mixture was then added in quadruplicate to 2.5×10^5 PBMCs. The culture continued for 3 days at which time cells were washed and V3
25 Loop-Toxin Chimeras 9952 resuspended in medium containing interleukin 2. Accumulation of p24 was detected by an ELISA.

Because the sera taken from one of the rabbits immunized with ntPE-V3MN26 neutralized virus in the MT4 assay at a dilution of 1:400, this serum was used to evaluate activity against the clinical isolates. A serum sample taken at 24 weeks
30 exhibited neutralizing activity against both a B and E subtype isolate (see Fig. 14). No neutralizing activity was seen with the pre-bleed sera from the same rabbit.

VI. ELICITATION OF IgA-MEDIATED IMMUNE RESPONSE

Mucosal inoculation by a PE-like chimeric immunogen containing 26 amino acids of the V3 loop of gp120 of HIV-1 induced both a humoral and cell-mediated immune response against HIV-1. A toxic version of this chimera was capable of killing a human intestinal cell line, Caco-2, grown as confluent monolayers. A non-lethal form of the chimera was administered to mice either subcutaneously or at vaginal, rectal, gastric or nasal mucosal surfaces. Subsequent boostings were performed at these various mucosal surfaces of by subcutaneous administration. Measurement of MNgp120-specific antibodies in serum and saliva samples demonstrated both IgA and IgG responses in every group of mucosal and subcutaneous administration. These results demonstrate that the PE-like chimeric immunogens of this invention can enter epithelial cells, be trafficked similarly to native toxin, transport across an intact epithelial barrier and induce the production of both IgA and IgG antibodies.

A. PE-Like Chimeric Immunogens

PE-like chimeras used in these experiments are described in Example I. The structural gene encoding native (toxic) PE was modified to delete the Ib region and provide a unique PstI site for the insertion of 26 amino acid V3 loop sequences. Non-toxic versions of PE-V3 loop chimeras were prepared which lacked the glutamic acid residue at position 553 ($\Delta E553$) and thus has no ADP-ribosylating activity. All PE-V3 loop chimeric proteins were expressed in *E coli* BL21(λ DE3) using the T7 promoter/T7 polymerase system. IPTG (1.0 mM for 90 min) was added to enhance protein expression. PE-V3 loop proteins were isolated from inclusion bodies and purified by successive rounds of anion exchange chromatography and a final gel filtration.

B. Cell-based Studies

A toxic version of the *Pseudomonas* exotoxin (PE) chimera containing 26 amino acids of the V3 loop of MN gp120 (tPE-MN26) was applied to the apical surface of confluent monolayers of polarized Caco-2 cells. Caco-2 cells were cultured and maintained as previously described (W. Rubas *et al.* (1996) "Flux measurements across Caco-2 monolayers might predict transport in human large intestinal tissue" *J. Pharm. Sci.* 85:165-169) on prewetted (PBS, 15 min. outside and then inside) collagen-coated polycarbonate filter supports (Snapwells™). Culture media was changed every other day

and confluent monolayers were used on day 25 post seeding and at passage 30-35. Toxic versions of PE and PE-V3 loop chimeras were added to the apical surface in culture media. After 24 h of continued incubation at 37 °C, Caco-2 monolayers were washed thrice with PBS to remove serum esterase activities and incubated with calcein AM and ethidium homodimer to determine live/dead cell ratio (LIVE/DEAD® Eukolight kit; Molecular Probes, Inc., Eugene, OR).

The chimera killed these intestinal epithelial cells with a potency similar to that of authentic PE (Fig. 8). Cell viability was measured as the ratio of live and dead cells.

A non-toxic ($\Delta 553$) chimera (ntPE-V3MN26) was used in all subsequent immunization studies to examine the ability of ntPE-V3MN26 to block the toxic actions of PE on Caco-2 monolayers (Fig. 8). Thus, the results of Figure 8 show that the incorporation of 26 amino acids of the V3 loop of MNgp120 (ntPE-V3MN26) in place of the endogenous Ib loop of PE does not alter the ability of the PE chimera to be taken up and processed by polarized, confluent epithelial cells. This ability of the PE-V3 loop chimera to be taken up and processed by epithelial cells is important against a pathogen such as HIV-1 which can infect and alter the function of human intestinal epithelial cells. D.M. Asmuth *et al.* (1994) "Physiological effects of HIV infection on human intestinal epithelial cells: an *in vitro* model for HIV enteropathy" *AIDS* 8:205-211.

C. Immunization Protocols

Female balb-c mice were obtained from Simonsen at 6 - 8 weeks of age and quarantined for 2 weeks prior to study. Animals were placed into one of 6 groups which were inoculated 3 times at two week intervals. The animals were maintained on ad lib food and water. Animal groups were immunized as follows: (1) oral, oral, oral; (2) vaginal, vaginal, vaginal; (3) rectal, rectal, rectal; (4) vaginal, oral, oral; (5) rectal, oral, oral; and (5) subcutaneous, subcutaneous, subcutaneous. Each oral inoculation used 40 μ g of PE-V3 loop chimera in 200 μ l of PBS containing 0.05% Tween 20, 1 mg/ml BSA and 0.2 M NaHCO₃ (pH = 8.1). All vaginal, rectal and subcutaneous inoculations contained 20 μ g PE-V3 loop chimera in 20 μ l of PBS containing 0.05% Tween 20.

D. Antibody Titers

Following intraperitoneal injection of 0.1 mg pilocarpine, mouse saliva (typically 50 μ l) was collected using polypropylene Pasteur pipettes and placed into polypropylene tubes. Serum samples (100 μ l) were obtained from periorbital bleeds using serum separator tubes. Collected serum and saliva samples were stored at -70 $^{\circ}$ C until analysis. A gp120-specific ELISA was performed using Costar 9018 E.I.A./R.I.A. plates coated with gp120. Following washing with PBST (PBS containing 0.05% Tween 20 and 0.01% thimerosal), plates were blocked with assay buffer (PBST containing 0.5% BSA). A subsequent washing was performed prior to serum or saliva sample introduction (100 μ l/well). Bound immunoglobulins were tagged using biotinylated whole goat antibodies which selectively recognized either mouse IgA or mouse IgG (Amersham). A mouse monoclonal antibody denoted 1F12 (Genentech, Inc.) was used as a positive control for IgG assays. No gp120-specific mouse IgA was available as a positive standard. ExtrAvidin[®] peroxidase conjugate (Sigma), 2,2'-azino-bis(2ethylbenzthiazoline-6-sulfonic acid (Sigma) and a phosphate-citrate buffer containing urea and hydrogen peroxide were used to quantitate bound antibody at 405 nm.

Non-toxic PE-V3MN26 was delivered to balb-c mice in combinations of oral gavage, application to the vaginal mucosa, application to the rectal mucosa or by subcutaneous injection. Serum and saliva samples were collected one, two and three months after the initial inoculation from each dosing group and analyzed by ELISA to determine IgG and IgA antibody titers specific for MN gp120. Pre-immune saliva and serum samples showed no significant background reaction in these gp120-specific ELISAs. Measurable quantities of gp120-specific IgG were observed in the sera of all dosing groups (Fig. 9). Although the IgG response observed was initially greatest in the subcutaneous group, all groups ultimately demonstrated strong serum IgG responses. Groups that were exposed orally to the ntPE-V3MN26 also appeared to obtain an IgG response faster than those groups exposed only at the vaginal or rectal mucosa. Compared to a mouse monoclonal IgG₁ which selectively recognizes the V3 loop of MNgp120, the highest measured levels in each of the groups of gp120-specific IgG were between 5-25 μ g/ml sera.

IgA antibodies appear to contribute to resistance against both strict mucosal pathogens and invasive agents which go on to cause systemic disease after mucosal colonization. R.I. Walker *et al.* (1994) "New strategies for using mucosal

vaccination to achieve more effective immunization" *Vaccine* 12:387-400. An ELISA was used to determine gp120-specific IgA levels in collected saliva samples as an index of mucosal antibody response. Since there is no MN gp120-specific monoclonal IgA available, values obtained by ELISA were only compared between groups and not characterized as absolute levels. Saliva samples from all 6 dosing groups contained gp120-specific IgA (Fig. 10). The strongest IgA response was observed in animals which received an initial vaginal dose and subsequent oral doses of PE-V3 loop chimera. It was interesting that animals which received only subcutaneous injections demonstrated IgA levels comparable to some of those observed in groups receiving only mucosal exposure of the chimera. This may be related to issues of the antibodies used in the IgA ELISA. Regardless, these results show that both mucosal and systemic immunity can be induced by mucosal immunization similar to that observed previously with oral immunization using pertussis toxin. M. J. Walker, *et al.* (1992) "Specific lung mucosal and systemic immune responses after oral immunization of mice with *Salmonella typhimurium* aro A, *Salmonella typhi* Ty21a, and invasive *Escherichia coli* expressing recombinant pertussis toxin S1 subunit" *Infect. Immun.* 60:4260.

HIV-1 subunit vaccines have been reported to only produce an IgG response following subcutaneous administration (M. B. Vasudevachari *et al.* (1992) "Envelope-specific antibodies in the saliva of individuals vaccinated with recombinant HIV-1 gp160" *J. Acquir. Immune Defic. Syndr.* 5:817-821) or both IgG and IgA following intramuscular injection (G. J. Gorse *et al.* (1996) "Salivary binding antibodies induced by human immunodeficiency virus type 1 recombinant gp120 vaccine" *Clin. Diagnostic Lab. Immunol.* 3:769-773.). Although those authors suggested that maximizing the production of mucosal antibodies will be important for an HIV-1 vaccine, it is unclear, however, if the IgA antibodies detected were secretory. It is likely that sIgA was the primary form of IgA in saliva samples and that dimeric IgA was the primary form in serum samples in those as well as the present studies. The IgA-binding reagent used presently was raised against serum IgA and thus may have provided a bias in IgA measurements. Thus the IgA levels measured in serum may only appear greater than saliva levels due to a lower affinity for sIgA than dimeric IgA. The IgA values given in the present study, therefore, are only presented on a relative scale.

A number of factors released by Th1 and Th2 cells have been shown to regulate IgA responses (J. R. McGhee *et al.* (1993) "New perspectives in mucosal

immunity with emphasis on vaccine development" *Seminars in Hematology*. 30:3-15). For example, in the presence of IL-5, IL-2 synergizes with TGF- β to augment IgA synthesis, leading to the prospect of pharmacologically manipulating the immune response. The form of antigen presentation, however, is dictated significantly by the fate of the immunogen. Epithelial cells at mucosal surfaces, which have the LRP receptor to bind and internalize ntPE-V3MN26, have been shown to express MHC class II proteins and class II can efficiently reach the surface of cells for antigen presentation from a lysosomal origin (V. G. Brachet *et al.* (1997) "Ii chain controls the transport of major histocompatibility complex class II molecules to and from lysosomes" *J. Cell Biol.* 137:51-65). Thus, ntPE-V3MN26 can be delivered by MHC class II structures onto the cell surface of epithelial cells. Alternatively, if the immunogen crosses the mucosal barrier and reaches a professional antigen presentation cell in the underlying lamina propria in an intact form, it should induce a Th2 response and result in a MHC class I-restricted antigen presentation.

VII. MEMORY RESPONSE ELICITED BY MUCOSAL ADMINISTRATION OF CHIMERIC IMMUNOGEN

Mucosal administration of ntPE-V3MN26 produced a significant memory response characterized by combination of serum IgG isotypes of both Th1 and Th2 pathways. Since the Th2 response has been proposed to be advantageous for neutralizing viruses and the cytotoxic immune responses associated with Th1 events may be required for effective immune responses against intracellular viruses (J.R. McGhee *et al.* (1994) *Reprod. Fertil. Dev.* 6:369-379), these results suggest that the mucosal immunization with ntPE-V3MN26 provided the types of responses desired for protection against HIV-1 infection (G.L. Ada *et al.* (1997) *AIDS Res. Hum. Retroviruses* 13:205-210).

A. Materials And Methods

1. Reagents

The structure and preparation of the ntPE-V3MN26 used in these studies is described herein. MNgp120 and the 1F12 monoclonal antibody recognizing the V3 loop of MNgp120 were prepared at Genentech, Inc. (South San Francisco, CA). Biotin-labeled goat antibodies raised against either mouse IgG or mouse IgA were purchased from Amersham Life Sciences (Arlington Heights, IL). Biotinylated rat antibodies

recognizing mouse IgG₁, IgG_{2a}, IgG_{2b}, IgG₃ and IgE were obtained from Pharmingen (San Diego, CA).

2. Immunization protocols and samples collection

Female BALB/c mice were obtained at 6-8 weeks of age and quarantined for 2 weeks prior to study and maintained throughout the study on ad lib food and water. Animals were randomly assigned to groups (n = 6) which received combinations of oral, vaginal, rectal or subcutaneous dosings. Oral inoculations were performed by oral gavage of 200 μ l of PBS containing 0.05% Tween 20, 1 mg/ml BSA, 0.2 M NaHCO₃ (final pH = 8.1) and 40 μ g of ntPE-V3MN26. Vaginal, rectal, and subcutaneous inoculations contained 20 μ g ntPE-V3MN26 in 20 μ l of PBS containing 0.05% Tween 20. Mouse saliva (typically 50-100 μ l) was collected over approximately 10 min using polypropylene Pasteur pipettes following hypersalivation induced by intraperitoneal injection of 0.1 mg pilocarpine per animal. Serum samples (100 μ l) were obtained from periorbital bleeds using serum separator tubes. Collected serum and saliva samples were stored at -70 °C until analysis.

In a separate study, mice were subcutaneously injected with 20 μ g ntPE-V3MN26 or 20 μ g ntPE and boosted at 2 and 7 weeks. One set of animals receiving ntPE-V3MN26 (n = 3) and the animals receiving ntPE (n = 2) were simultaneously dosed with 40 μ l of Freund's complete adjuvant initially and 40 μ l of Freund's incomplete adjuvant at weeks 2 and 7. A set of animals (n = 3) dosed with 20 μ g of ntPE-V3MN26 formulated in 40 μ l of normal saline served as a control. Serum samples (100 μ l) were obtained on a weekly basis and stored as described above.

3. Measurement of antibody responses

Anti-gp120-specific antibodies were measured by enzyme-linked immunosorbent assay (ELISA). Briefly, Costar 9018 E.I.A./R.I.A. 96-well plates were coated with 1 μ g/well of MNgp120, washed thrice with PBS containing 0.05% Tween 20 (v/v) and then blocked overnight at 4° C with PBS containing 1% BSA. After washing with PBS/Tween 20, plates were incubated with dilutions of serum or saliva samples (diluted with PBS/Tween 20 containing 0.1% BSA). The plates were incubated for 2 h at room temperature with gentle agitation, then washed thrice with PBS/Tween 20 and incubated with a biotin-conjugated goat anti-mouse IgA or IgG or, to determine IgG

subclass or IgE responses, with biotin-conjugated rat anti-mouse IgG1, IgG2a, IgG2b, IgG3, or IgE for 1 h using the same incubation conditions. After washing with PBS/Tween 20, horseradish peroxidase-conjugated streptavidin was added. Bound antibodies were visualized by ExtrAvidin[®] peroxidase conjugate (Sigma), 2,2'-azino-bis(2ethylbenzthiazoline-6-sulfonic acid (Sigma) and a phosphate-citrate buffer containing urea and hydrogen peroxide were used to quantitate bound antibody at 405 nm.

B. Results

1. IgA antibody responses to ntPE-V3MN26

Animals were inoculated (n = 6/group) by a variety of routes with ntPE-V3MN26 followed by 2 boosts on days 14 and 21 and then at month 16. Animals received ntPE-V3MN26 either orally (PO), vaginally (V), rectally (R), vaginally and orally (V/PO), rectally and orally (R/PO), or subcutaneously (SC). Saliva samples collected at 30, 60 and 90 days and then again at 16.5 months were analyzed for antigen-specific IgA (Fig. 11). Without an anti-V3 loop IgA antibody to standardize the assays, responses were normalized against one strongly positive sample. Values were reported on an arbitrary scale of antigen-specific IgA units. All dosing groups demonstrated comparable salivary IgA responses at 30 and 60 days. By 90 days, the strongest salivary IgA response was observed in the group which received an initial vaginal dose and subsequent oral boosts. At 16.5 months the all oral, all vaginal and all rectal groups showed the greatest levels of antigen-specific salivary IgA. Responses of the combined mucosal inoculation groups (vaginal/oral and rectal/oral) were comparable to those observed in the group dosed subcutaneously.

To insure that these salivary IgA responses reflected antigen-specific binding and not a non-specific binding to salivary components, pre-immune saliva samples were evaluated and a study was performed in which a mixture of V3 loop peptide and ntPE was administered to mice. The studies showed that undiluted pre-immune saliva samples did not demonstrate a measurable background in the ELISA format. Also, animals dosed simultaneously with ntPE and an unconjugated V3 loop constrained by a disulfide bond did not have measurable MNgp120-specific IgA levels. These results indicate that there was little or no non-specific cross-reactivity in the ELISA.

No detectable antigen-specific serum IgA responses were observed in any of the dosing groups at the 1, 2 or 3 month sampling times. However, at 16.5 months, sera collected from all groups demonstrated antigen-specific IgA (Table 1). It is possible that the ability to detect serum IgA at this time may have been due to a heightened total immune response rather than a specific stimulation. Interestingly, the relative serum IgA levels did not correlate with salivary IgA levels. For example, rectal/oral combination inoculations yielded one of the weaker memory salivary IgA responses but the strongest memory serum IgA response (Table 1, Fig. 11). The all oral, all vaginal or all rectal groups, which provided the greatest salivary IgA responses at 16.5 months had some of the weakest serum IgA responses at this time. Unlike mucosal administration of ntPE-V3MN26 where opposing levels in saliva and serum were the norm, subcutaneous inoculations of ntPE-V3MN26 produced a moderate IgA response in both the saliva and serum of mice (Table 1, Fig. 11). Whatever the stimulus of IgA production, the antigen-specific serum IgA levels were transient. At the 22 month sampling, just two animals of the rectal/oral group represented the only positives for measurable serum IgA recognizing MNgp120. No other groups, even the subcutaneous injection group, showed any detectable serum IgA levels at this time point.

TABLE 1. Immunization with ntPE-V3MN26 stimulates the production of antigen-specific serum IgA and salivary IgG in Mice

Immunization schedule ^a	Serum IgA ^b (arbitrary units)	Salivary IgG ^c (μ g/ml)
PO/PO/PO/PO	0.233 \pm 0.074	10.9 \pm 2.2
V/V/V/V	0.172 \pm 0.061	9.52 \pm 1.6
R/R/R/R	0.178 \pm 0.042	9.93 \pm 1.7
V/PO/PO/PO	0.160 \pm 0.021	9.90 \pm 1.3
R/PO/PO/PO	0.450 \pm 0.128	11.0 \pm 0.49
SC/SC/SC/SC	0.273 \pm 0.078	7.1 \pm 0.63

^a Immunizations were performed at days 0, 14, 21 and at month 16 to animals either orally (PO) vaginally (V), rectally (R) or subcutaneously (SC).

^b MNgp120-specific IgA levels were measured by ELISA at 16.5 months and normalized against a single sample standard and reported in arbitrary units.

^c MNgp120-specific IgG levels were measured by ELISA at 16.5 months and calibrated against a mouse monoclonal antibody (1F12) which recognizes the V3 loop of the protein.

2. IgG antibody responses to ntPE-V3MN26

Serum and salivary antigen-specific IgG responses, measured by ELISA, were standardized using a mouse monoclonal antibody (1F12) which recognizes the V3 loop of MNgp120. The assay was linear over the range of 0.05 - 2.5 μ g for 1F12 and pre-immune sera and salivas were negative in the ELISA format. Although the IgG response produced by an initial inoculation followed by two boosts was ultimately greatest in the subcutaneous injection group, all mucosal inoculation groups demonstrated strong serum IgG responses at 30, 60 and 90 days (Fig. 12). Two weeks after an ntPE-V3MN26 boost at month 16 the subcutaneous injection group had the highest serum IgG memory response. All mucosal groups also showed strong memory responses at this time (Fig. 12). However, by month 22 antigen-specific serum IgG titers had decreased in all groups.

3. Comparison of serum and saliva IgG and IgA levels

Previous studies have suggested that serum IgG can transudate onto mucosal surfaces, possibly providing some form of immune protection. M.B. Vasudevachari et al. (1992) *J. Acquir. Immune Defic. Syndr.* 5:817-821. Others have not been able to demonstrate such a transudative event. E.-L. Johansson et al. (1998) *Infect. Immun.* 66:514-520. In these studies, antigen-specific IgG was not observed in saliva samples at months 1, 2 and 3 but rose to detectable levels following a boost at month 16 (Table 1). All mucosally dosed animal groups had comparable salivary IgG responses at this time which were greater than that observed for the animals receiving subcutaneous ntPE-V3MN26 (Table 1). This lack of correlation between relative serum and saliva levels of antigen-specific IgG (Fig. 12, Table 1) suggests a separation of the serum and salivary IgG pools resulting from this memory response. Thus, it appears that the IgG present in saliva in the studies may have resulted, to a significant extent, from local antibody production rather than a "spill-over" from circulating serum antibodies.

4. Serum IgG isotype responses to ntPE-V3MN26

In mice, induction of a Th1 response typically leads to the production of IgG2a and IgG3 by B cells while a Th2 response results in IgG1 and possibly IgE production. A.K. Abbas et al. (1996) *Nature* 383:787-793. The development of either a Th1 or Th2 response is driven by specific cytokines such as interferon- γ and IL-4. Introduction of ntPE-V3MN26 either systemically through subcutaneous injection or via application at oral, vaginal or rectal tissues led to the development of an antigen-specific serum IgG response. The IgG isotype population of these sera samples was investigated and it was found that the MNgp120-specific response was dominated (~55%) by IgG1. Lesser and comparable amounts of antigen-specific IgG2a (~20%) and IgG2b (~20%) were found along with low amounts (~5%) of IgG3. No antigen-specific IgE was detected. These results suggest that subcutaneous administration of ntPE-V3MN26 induces both Th1 and Th2 responses in BALB/c mice with the Th2 phenotype dominating.

VIII. EVALUATION OF ntPE-V3MN26 AS AN ADJUVANT

Adjuvants can act to facilitate the presentation of an antigen and/or activate the immune response at the site of inoculation. F.R. Vogel et al. (1995) *A compendium of vaccine adjuvants and excipients*, p. 141-228. In M. F. Powell, and M. J. Newman (ed.), *VACCINE DESIGN: THE SUBUNIT AND ADJUVANT APPROACH*, vol. 6. Plenum Press, New York. Recognized as one of the most potent adjuvants available, Freund's adjuvant is a mixture of mineral oil, surfactant and *Mycobacterium tuberculosis*. A study to assess the efficiency of serum IgG induction by ntPE-V3MN26 was performed by injecting mice subcutaneously with ntPE-V3MN26 and Freund's complete adjuvant initially, boosting with ntPE-V3MN26 and incomplete adjuvant after 14 and 49 days, and then comparing IgG serum responses to those of animals receiving ntPE-V3MN26 without Freund's adjuvant (Fig. 13). Animals receiving the same subcutaneous dosing regime of ntPE-V3MN26 with normal saline instead of Freund's adjuvant exhibited approximately one-third the antigen-specific immune response that observed in animals receiving this chimera along with Freund's adjuvant. The level of response to ntPE-V3MN26 over this time frame was similar to that observed in the subcutaneous injection group graphed in Fig. 12 at months 1, 2 and 3, suggesting a fairly consistent outcome for this form of chimera delivery. A control where the Freund's adjuvant regimen was

injected along with a non-toxic PE which lacked the V3 loop of MNgp120 demonstrated the specificity of the immune response being measured (Fig. 13).

5 The present invention provides *Pseudomonas* exotoxin A-like chimeric immunogens and methods of evoking an immune response. While specific examples have been provided, the above description is illustrative and not restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification. The scope of the invention should, therefore, be determined not with reference to the above description, but instead should be determined with reference to the
10 appended claims along with their full scope of equivalents.

All publications and patent documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication or patent document were so individually denoted. By their citation of various references in this document Applicants do not admit that any particular
15 reference is "prior art" to their invention.

WHAT IS CLAIMED IS:

1 1. A non-toxic *Pseudomonas* exotoxin A-like ("PE-like") chimeric
2 immunogen comprising: (1) a cell recognition domain of between 10 and 1500 amino
3 acids that binds to a cell surface receptor; (2) a translocation domain comprising an
4 amino acid sequence substantially identical to a sequence of PE domain II sufficient to
5 effect translocation to a cell cytosol; (3) a non-native epitope domain comprising an
6 amino acid sequence of between 5 and 1500 amino acids that encodes a non-native
7 epitope; and (4) an amino acid sequence encoding an endoplasmic reticulum ("ER")
8 retention domain that comprises an ER retention sequence.

1 2. The immunogen of claim 1 having the amino acid sequence of PE
2 Δ E553 except that the sequence of domain Ib of PE Δ E553 comprises the non-native
3 epitope between two cysteine residues of domain Ib.

1 3. The immunogen of claim 1 wherein the cell recognition domain is
2 domain Ia of PE.

1 4. The immunogen of claim 1 wherein cell recognition domain binds
2 to α 2-macroglobulin receptor (" α 2-MR"), epidermal growth factor ("EGF") receptor; the
3 IL-2 receptor; the IL-6 receptor; HIV-infected cells; a chemokine receptor; a leukocyte
4 cell surface receptor; a ligand for the IgA receptor; or an antibody or antibody fragment
5 directed to a receptor.

1 5. The immunogen of claim 1 wherein cell recognition domain
2 comprises amino acid sequences of a growth factor or an antibody.

1 6. The immunogen of claim 1 wherein cell recognition domain is
2 comprised within the ER retention domain.

1 7. The immunogen of claim 1 wherein the translocation domain
2 comprises amino acids 280 to 364 of domain II of PE.

1 8. The immunogen of claim 1 wherein the translocation domain is
2 domain II of PE.

1 9. The immunogen of claim 1 wherein the non-native epitope domain
2 comprises a cysteine-cysteine loop that comprises the non-native epitope.

1 10. The immunogen of claim 1 wherein the non-native epitope domain
2 comprises an amino acid sequence encoding a non-native epitope inserted between two
3 cysteine residues of domain Ib of PE.

1 11. The immunogen of claim 1 wherein the non-native epitope domain
2 comprises an amino acid sequence selected from CTRPNYNKRK RIHIGPGRAF
3 YTTKNIIGTI RQAHC (SEQ ID NO:3) or CTRPSNNTRT SITIGPGQVF YRTGDIIGDI
4 RKAYC (SEQ ID NO:4).

1 12. The immunogen of claim 1 wherein the ER retention domain is
2 domain III of PE comprising the mutation Δ E553.

1 13. The immunogen of claim 1 wherein the ER retention sequence
2 comprises REDLK (SEQ ID NO:11), REDL (SEQ ID NO:12) or KDEL (SEQ ID
3 NO:13).

1 14. The immunogen of claim 1 which is ntPE-V3MN14 or ntPE-
2 V3MN26.

1 15. The immunogen of claim 1 wherein the non-native epitope is an
2 epitope from a viral, bacterial or parasitic protozoan pathogen.

1 16. The immunogen of claim 9 wherein the non-native epitope is an
2 epitope of a V3 loop of gp120 of HIV-1.

1 17. The immunogen of claim 9 wherein the non-native epitope is an
2 epitope of a principal neutralizing loop of a retrovirus.

1 18. The immunogen of claim 9 wherein the non-native epitope is an
2 epitope of a major neutralizing loop of HIV-2 or a V3 loop of gp120 of HIV-1 of at least
3 8 amino acids including a V3 loop apex.

1 19. A recombinant polynucleotide comprising a nucleotide sequence
2 encoding a non-toxic *Pseudomonas* exotoxin A-like ("PE-like") chimeric immunogen, the
3 PE-like chimeric immunogen comprising: (1) a cell recognition domain of between 10
4 and 1500 amino acids that binds to a cell surface receptor; (2) a translocation domain
5 comprising an amino acid sequence substantially identical to a sequence of PE domain II
6 sufficient to effect translocation to a cell cytosol; (3) a non-native epitope domain
7 comprising an amino acid sequence of between 5 and 1500 amino acids that encodes a
8 non-native epitope; and (4) an amino acid sequence encoding an endoplasmic reticulum
9 ("ER") retention domain that comprises an ER retention sequence.

1 20. The recombinant polynucleotide of claim 19 which is an expression
2 vector further comprising an expression control sequence operatively linked to the
3 nucleotide sequence.

1 21. The recombinant polynucleotide of claim 19 having the amino acid
2 sequence of PE wherein domain Ib of PE further comprises the non-native epitope
3 between two cysteine residues of domain Ib.

1 22. A recombinant non-toxic *Pseudomonas* exotoxin A-like ("PE-like")
2 chimeric immunogen cloning platform comprising a nucleotide sequence encoding: (1) a
3 cell recognition domain of between 10 and 1500 amino acids that binds to a cell surface
4 receptor; (2) a translocation domain comprising an amino acid sequence substantially
5 identical to a sequence of PE domain II sufficient to effect translocation to a cell cytosol;
6 (3) an amino acid sequence encoding an endoplasmic reticulum ("ER") retention domain
7 that comprises an ER retention sequence and (4) a splicing site between the sequence
8 encoding the translocation domain and the sequence encoding the ER retention domain.

1 23. The recombinant cloning platform of claim 22 which is an
2 expression vector further comprising an expression control sequence operatively linked to
3 the nucleotide sequence.

1 24. A method of producing antibodies against a non-native epitope,
2 wherein the non-native epitope naturally exists within a cysteine-cysteine loop comprising
3 the step of inoculating an animal with a non-toxic *Pseudomonas* exotoxin A-like ("PE-
4 like") chimeric immunogen, the PE-like chimeric immunogen comprising: (1) a cell
5 recognition domain of between 10 and 1500 amino acids that binds to a cell surface
6 receptor; (2) a translocation domain comprising an amino acid sequence substantially
7 identical to a sequence of PE domain II sufficient to effect translocation to a cell cytosol;
8 (3) a non-native epitope domain comprising a cysteine-cysteine loop that contains within
9 the loop an amino acid sequence of between 5 and 1500 amino acids that encodes a non-
10 native epitope; and (4) an amino acid sequence encoding an endoplasmic reticulum
11 ("ER") retention domain that comprises an ER retention sequence.

1 25. The method of claim 24 wherein the cysteine-cysteine loop
2 comprises no more than about 30 amino acids.

1 26. The method of claim 24 wherein the non-native epitope is an
2 epitope of the V3 domain of HIV-1.

1 27. A vaccine comprising at least one non-toxic *Pseudomonas* exotoxin
2 A-like ("PE-like") chimeric immunogen, the PE-like chimeric immunogen comprising:
3 (1) a cell recognition domain of between 10 and 1500 amino acids that binds to a cell
4 surface receptor; (2) a translocation domain comprising an amino acid sequence
5 substantially identical to a sequence of PE domain II sufficient to effect translocation to a
6 cell cytosol; (3) a non-native epitope domain comprising an amino acid sequence of
7 between 5 and 1500 amino acids that encodes a non-native epitope; and (4) an amino
8 acid sequence encoding an endoplasmic reticulum ("ER") retention domain that
9 comprises an ER retention sequence.

28. The vaccine of claim 27 comprising a plurality of PE-like chimeric immunogens, each immunogen having a different non-native epitope.

29. The vaccine of claim 27 further comprising a pharmaceutically acceptable carrier.

30. The vaccine of claim 27 in the form of an immunization dose wherein the immunogen is present in an amount effective to elicit in a human subject an immune response against the non-native epitope.

31. The vaccine of claim 28 wherein the different non-native epitopes are epitopes of different strains of the same pathogen.

32. The vaccine of claim 31 wherein the non-native epitope is an epitope of the V3 loop of HIV-1 and the different strains of the same pathogen are HIV-1 MN and HIV-1 Thai-E.

33. A method of eliciting an immune response against a non-native epitope in a subject, the method comprising the step of administering to the subject a vaccine comprising at least one non-toxic *Pseudomonas* exotoxin A-like ("PE-like") chimeric immunogen, the PE-like chimeric immunogen comprising: (1) a cell recognition domain of between 10 and 1500 amino acids that binds to a cell surface receptor; (2) a translocation domain comprising an amino acid sequence substantially identical to a sequence of PE domain II sufficient to effect translocation to a cell cytosol; (3) a non-native epitope domain comprising an amino acid sequence of between 5 and 1500 amino acids that encodes a non-native epitope; and (4) an amino acid sequence encoding an endoplasmic reticulum ("ER") retention domain that comprises an ER retention sequence.

34. The method of claim 33 wherein the non-native epitope comprises a binding motif for an MHC Class II molecule of the subject and the immune response elicited is an MHC Class-II dependent cell-mediated immune response.

1 35. The method of claim 33 wherein the non-native epitope comprises a
2 binding motif for an MHC Class I molecule of the subject and the immune response
3 elicited is an MHC Class-I dependent cell-mediated immune response.

1 36. The method of claim 33 wherein the non-native epitope is an
2 epitope of the V3 domain of HIV-1.

1 37. The method of claim 33 wherein the vaccine is administered as a
2 prophylactic treatment against a disease mediated by an agent bearing the non-native
3 epitope.

1 38. The method of claim 33 wherein the vaccine is administered as a
2 therapeutic treatment against a disease mediated by an agent bearing the non-native
3 epitope.

1 39. A polynucleotide vaccine comprising at least one recombinant
2 polynucleotide comprising a nucleotide sequence encoding a non-toxic *Pseudomonas*
3 exotoxin A-like ("PE-like") chimeric immunogen, the PE-like chimeric immunogen
4 comprising: (1) a cell recognition domain of between 10 and 1500 amino acids that binds
5 to a cell surface receptor; (2) a translocation domain comprising an amino acid sequence
6 substantially identical to a sequence of PE domain II sufficient to effect translocation to a
7 cell cytosol; (3) a non-native epitope domain comprising an amino acid sequence of
8 between 5 and 1500 amino acids that encodes a non-native epitope; and (4) an amino
9 acid sequence encoding an endoplasmic reticulum ("ER") retention domain that
10 comprises an ER retention sequence.

1 40. A method of eliciting an immune response against a non-native
2 epitope in a subject, the method comprising the step of administering to the subject a
3 polynucleotide vaccine comprising at least one recombinant polynucleotide comprising a
4 nucleotide sequence encoding a non-toxic *Pseudomonas* exotoxin A-like ("PE-like")
5 chimeric immunogen, the PE-like chimeric immunogen comprising: (1) a cell recognition
6 domain of between 10 and 1500 amino acids that binds to a cell surface receptor; (2) a
7 translocation domain comprising an amino acid sequence substantially identical to a

sequence of PE domain II sufficient to effect translocation to a cell cytosol; (3) a non-native epitope domain comprising an amino acid sequence of between 5 and 1500 amino acids that encodes a non-native epitope; and (4) an amino acid sequence encoding an endoplasmic reticulum ("ER") retention domain that comprises an ER retention sequence.

41. The method of claim 40 wherein the recombinant polynucleotide is an expression vector comprising an expression control sequence operatively linked to the nucleotide sequence.

42. The method of claim 40 wherein the nucleotide sequence further encodes a mammalian secretory sequence attached to the amino terminus of the immunogen.

43. A method of eliciting an immune response against a non-native epitope in a subject, the method comprising the steps of transfecting cells with a recombinant polynucleotide comprising a nucleotide sequence encoding a non-toxic *Pseudomonas* exotoxin A-like ("PE-like") chimeric immunogen, the PE-like chimeric immunogen comprising: (1) a cell recognition domain of between 10 and 1500 amino acids that binds to a cell surface receptor; (2) a translocation domain comprising an amino acid sequence substantially identical to a sequence of PE domain II sufficient to effect translocation to a cell cytosol; (3) a non-native epitope domain comprising an amino acid sequence of between 5 and 1500 amino acids that encodes a non-native epitope; and (4) an amino acid sequence encoding an endoplasmic reticulum ("ER") retention domain that comprises an ER retention sequence, and administering the cells to the subject.

PSEUDOMONAS EXOTOXIN A-LIKE CHIMERIC IMMUNOGENS


ABSTRACT OF THE DISCLOSURE

This invention provides *Pseudomonas* exotoxin A-like chimeric immunogens that include a non-native epitope in the Ib domain of *Pseudomonas* exotoxin. Methods of eliciting an immune response using these immunogens also are provided.


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195

1/14

1 A Pseudomonas exotoxin



1 B V3-loop toxin chimera




1 C

Protein	Ib Loop Region Amino Acid Sequence	Molecular Mass
wild-type PE	-GAANADVVS LTCPVAAGECAGPAD-	67,122 Da
ntPE-V3MN14	-GAANLHCGIHIGPGRAFYTTKCMQGPAD-	67,729 Da
ntPE-V3MN26	-GAANLHCNYNKRKRIHIGPGRAFYT TKNIIGTICMQGPAD-	68,937 Da
ntPE-V3ThE26	-GAANLHCNNTRTSITIGPGQVFYRT GDIIGDDICMQGPAD-	68,700 Da
ntPE-FP16	-GAANLQCRLEEKKGNYVVT DHR LCLQGPAD-	67,862 Da

Figure 1.


09/462682.042800

2/14



SDS-PAGE

FIG. 2A.




IF12/MN

Western blot Analysis


FIG. 2B.



IB2/Th-E



IF12/MN



IB2/Th-E


Antibody-mediated capture of proteins

FIG. 2C.


SUBSTITUTE SHEET (RULE 26)

3/14

3A



3B



3C

Sample	% α -helix*	% β -sheet	% β -turns	% other
wild-type PE	21	48	12	20
ntPE-V3MN14	26	46	10	19
ntPE-V3MN26	18	50	12	20

Figure 3.
 SUBSTITUTE SHEET (RULE 26)

4/14

Cytotoxic activity for A431 Cells




Figure 4.

5/14




FIG. 5A.




FIG. 5B.

6/14




Figure 6.

PCT/US 98/14341
RO/US 30 OCT 1998
09/462682

* FURIN CLEAVAGE SITE (279-280)

 = HATCHED AREA NOT ESSENTIAL FOR TRANSLOCATION

Figure 7.

8/14




Figure 8.

9/14





Figure 9.

10/14

IgA Levels in Pooled Salivas



Normalized gp120-Specific ELISA Values

Figure 10.

Salivary IgA Response Induced by ntPE-MN26




Figure 11.

11/14

SUBSTITUTE SHEET (RULE 26)

PCT/US 98/14341
RO/US 30 OCT 1998
09/462682

09/52682 09/2800

Serum IgG Response to Subcutaneous Injection of nPE-MN26




Figure 12.

09452632, 042800

12/14

PCT/US 98/114341
RO/US 30 OCT 1998
09/462682

Serum IgG Response Induced by ntPE-MN26





Figure 13.

V3 Loop-Toxin Chimeras
Growth of HIV-1/RVL05



Growth of HIV-1/Th92009




Figure 14.

14/14

PCT/US 98/14341
RO/US 30 OCT 1998
09/462682

09462682 043800

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: PSEUDOMONAS EXOTOXIN A-LIKE CHIMERIC IMMUNOGENS the specification of which is attached hereto or X was filed on January 10, 2000 as Application No. 09/462,682 and was amended on (if applicable).

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56. I claim foreign priority benefits under Title 35, United States Code, Section 119 of any foreign applications(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Country	Application No.	Date of Filing	Priority Claimed Under 35 USC 119
PCT	PCT/US98/14341	July 10, 1998	Yes No
			Yes No

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

Application No.	Filing Date
60/052,375	July 11, 1997

I claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Date of Filing	Status
		Patented Pending Abandoned
		Patented Pending Abandoned
		Patented Pending Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) of the Government of the United States of America as represented by the Secretary of the Department of Health and Human Services, Office of Technology Transfer, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Robert Benson, Reg. No. 33,612
 Steven M. Ferguson, Reg. No. 38,448
 James C. Haight, Reg. No. 25,588
 John P. Kim, Reg. No. 38,514

37

Susan S. Rucker, Reg. No. 35,762
 David R. Sadowski, Reg. No. 32,808
 Jack Spiegel, Reg. No. 34,477




Send Correspondence to Appointed Associate Attorney or Agent Address: TOWNSEND and TOWNSEND and CREW, LLP Two Embarcadero Center, 8th Floor San Francisco, CA 94111-3834	Direct Telephone Calls and Facsimiles to Appointed Associate Attorney or Agent: Name: John Storella Reg. No. 32,944 Telephone: 415-576-0200 Fax: 415-576-0300
--	---

1-00

Full Name of Inventor 1	Last Name <u>FITZGERALD</u>	First Name <u>DAVID</u>	Middle Name or Initial <u>J.</u>	
Residence & Citizenship	City <u>Rockville</u>	State/Foreign Country Maryland <u>MD</u>	Country of Citizenship United States	
Post Office Address	Post Office Address 1202 Azalea Drive	City Rockville	State/Country MD	Zip Code 20850

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Title 18 of the United States Code, Section 1001 and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1  David J. Fitzgerald		
Date <u>4/15/00</u>		

SF 1083566 v1

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

FitzGerald

Serial No.: 09/462,682

Filed: January 10, 2000

For: PSEUDOMONAS EXOTOXIN A-
LIKE CHIMERIC IMMUNOGENSAssistant Commissioner for Patents
Washington, D.C. 20231

Examiner: Unassigned

Art Unit: Unassigned

APPOINTMENT OF ASSOCIATE
ATTORNEYS AND AGENTS BY
PRINCIPAL ATTORNEY OR AGENT
UNDER 37 CFR §1.34(b) AND MPEP
402.02

Sir:

Under 37 CFR §1.34(b) and MPEP 402.02, the undersigned principal attorney or agent of record hereby appoints the following representatives from the law firm of TOWNSEND and TOWNSEND and CREW, LLP, Two Embarcadero Center, 8th Floor, San Francisco, California 94111-3834, telephone: (415) 576-0200, as associate attorneys and/or patent agents in this application with full powers to prosecute this case and to transact all business in the Patent and Trademark Office connected therewith:

Philip H. Albert, Reg. No. 35,819
Hector A. Alicea, Reg. No. 40,891
Randolph T. Apple, Reg. No. 36,429
Scott L. Ausenius, Reg. No. 42,271
Krishna G Banerjee, Reg. No. 43,317
Kevin L. Bastian, Reg. No. 34,774
Guy Chambers, Reg. No. 30,617
Kathleen L. Choi, Reg. No. 43,433
Karen B. Dow, Reg. No. 29,684
James F. Hann, Reg. No. 29,719
M. Henry Helmes, Reg. No. 28,219
James M. Heslin, Reg. No. 29,541
Matthew E. Hirsch, Reg. No. Not Yet Assigned
Laurence J. Hyman, Reg. No. 35,551
William Michael Hynes, Reg. No. 24,168
Jeffrey J. King, Reg. No. 38,515

Charlie E. Krueger, Reg. No. 30,077
Joe Liebeschuetz, Reg. No. 37,505
Jeffrey S. Mann, Reg. No. 42,837
Annette S. Parent, Reg. No. 42,058
Steven W. Parmelee, Reg. No. 31,990
Brian W. Poor, Reg. No. 32,928
Timothy L. Smith, Reg. No. 35,367
William M. Smith, Reg. No. 30,223
Andrew T. Serafini, Reg. No. 41,303
Joseph R. Snyder, Reg. No. 39,381
John R. Storella, Reg. No. 32,944
Eugenia Garrett-Wackowski, Reg. No. 37,330
Ellen Lauver Weber, Reg. No. 32,762
Kenneth A. Weber, Reg. No. 31,667
Michael E. Woods, Reg. No. 33,465

Dated: 4/17/00

Respectfully submitted,

Office of Technology Transfer
National Institutes of Health
6011 Executive Boulevard, Suite 325
Rockville, Maryland 20852
Telephone: (301) 496-7056

United States Department of Health and
Human Services

By Robert BensonReg. No. 33,612

SEQUENCE LISTING

The nucleotide sequence (SEQ ID NO:1) and deduced amino acid sequence (SEQ ID NO:2) of *Pseudomonas* exotoxin A are:

5	GCC	GAA	GAA	GCT	TTC	GAC	CTC	TGG	AAC	GAA	TGC	GCC	AAA	GCC	TGC	GTG	48
	Ala	Glu	Glu	Ala	Phe	Asp	Leu	Trp	Asn	Glu	Cys	Ala	Lys	Ala	Cys	Val	
	1				5					10				15			
10	CTC	GAC	CTC	AAG	GAC	GGC	GTG	CGT	TCC	AGC	CGC	ATG	AGC	GTC	GAC	CCG	96
	Leu	Asp	Leu	Lys	Asp	Gly	Val	Arg	Ser	Ser	Arg	Met	Ser	Val	Asp	Pro	
				20					25					30			
15	GCC	ATC	GCC	GAC	ACC	AAC	GGC	CAG	GGC	GTG	CTG	CAC	TAC	TCC	ATG	GTC	144
	Ala	Ile	Ala	Asp	Thr	Asn	Gly	Gln	Gly	Val	Leu	His	Tyr	Ser	Met	Val	
			35					40					45				
20	CTG	GAG	GGC	GGC	AAC	GAC	GCG	CTC	AAG	CTG	GCC	ATC	GAC	AAC	GCC	CTC	192
	Leu	Glu	Gly	Gly	Asn	Asp	Ala	Leu	Lys	Leu	Ala	Ile	Asp	Asn	Ala	Leu	
		50					55					60					
25	AGC	ATC	ACC	AGC	GAC	GGC	CTG	ACC	ATC	CGC	CTC	GAA	GGC	GGC	GTC	GAG	240
	Ser	Ile	Thr	Ser	Asp	Gly	Leu	Thr	Ile	Arg	Leu	Glu	Gly	Gly	Val	Glu	
	65					70				75						80	
30	CCG	AAC	AAG	CCG	GTG	CGC	TAC	AGC	TAC	ACG	CGC	CAG	CGC	CGC	GGC	AGT	288
	Pro	Asn	Lys	Pro	Val	Arg	Tyr	Ser	Tyr	Thr	Arg	Gln	Ala	Arg	Gly	Ser	
					85					90				95			
35	TGG	TCG	CTG	AAC	TGG	CTG	GTA	CCG	ATC	GGC	CAC	GAG	AAG	CCC	TCG	AAC	336
	Trp	Ser	Leu	Asn	Trp	Leu	Val	Pro	Ile	Gly	His	Glu	Lys	Pro	Ser	Asn	
				100				105						110			
40	ATC	AAG	GTG	TTC	ATC	CAC	GAA	CTG	AAC	GCC	GGC	AAC	CAG	CTC	AGC	CAC	384
	Ile	Lys	Val	Phe	Ile	His	Glu	Leu	Asn	Ala	Gly	Asn	Gln	Leu	Ser	His	
			115					120					125				
45	ATG	TCG	CCG	ATC	TAC	ACC	ATC	GAG	ATG	GGC	GAC	GAG	TTG	CTG	GCG	AAG	432
	Met	Ser	Pro	Ile	Tyr	Thr	Ile	Glu	Met	Gly	Asp	Glu	Leu	Leu	Ala	Lys	
			130				135					140					
50	CTG	GCG	CGC	GAT	GCC	ACC	TTC	TTC	GTC	AGG	GCG	CAC	GAG	AGC	AAC	GAG	480
	Leu	Ala	Arg	Asp	Ala	Thr	Phe	Phe	Val	Arg	Ala	His	Glu	Ser	Asn	Glu	
					150						155					160	
55	ATG	CAG	CCG	ACG	CTC	GCC	ATC	AGC	CAT	GCC	GGG	GTC	AGC	GTG	GTC	ATG	528
	Met	Gln	Pro	Thr	Leu	Ala	Ile	Ser	His	Ala	Gly	Val	Ser	Val	Val	Met	
					165					170				175			
60	GCC	CAG	ACC	CAG	CCG	CGC	CGG	GAA	AAG	CGC	TGG	AGC	GAA	TGG	GCC	AGC	576
	Ala	Gln	Thr	Gln	Pro	Arg	Arg	Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser	
				180					185					190			
65	GGC	AAG	GTG	TTG	TGC	CTG	CTC	GAC	CCG	CTG	GAC	GGG	GTC	TAC	AAC	TAC	624
	Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr	
			195					200					205				
70	CTC	GCC	CAG	CAA	CGC	TGC	AAC	CTC	GAC	GAT	ACC	TGG	GAA	GGC	AAG	ATC	672
	Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile	
			210				215					220					
75	TAC	CGG	GTG	CTC	GCC	GGC	AAC	CCG	GCG	AAG	CAT	GAC	CTG	GAC	ATC	AAA	720
	Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys	
						230					235					240	

5	CCC Pro	ACG Thr	GTC Val	ATC Ile	AGT Ser 245	CAT His	CGC Arg	CTG Leu	CAC His 250	TTT Phe 250	CCC Pro	GAG Glu	GGC Gly	GGC Gly	AGC Ser 255	CTG Leu	768
	GCC Ala	GCG Ala	CTG Leu	ACC Thr 260	GCG Ala	CAC His	CAG Gln	GCT Ala 265	TGC Cys	CAC His	CTG Leu	CCG Pro	CTG Leu	GAG Glu 270	ACT Thr	TTC Phe	816
10	ACC Thr	CGT Arg	CAT His 275	CGC Arg	CAG Gln	CCG Pro	CGC Arg	GGC Gly 280	TGG Trp	GAA Glu	CAA Gln	CTG Leu	GAG Glu 285	CAG Gln	TGC Cys	GGC Gly	864
15	TAT Tyr	CCG Pro 290	GTG Val	CAG Gln	CGG Arg	CTG Leu	GTC Val 295	GCC Ala	CTC Leu	TAC Tyr	CTG Leu	GCG Ala 300	GCG Ala	CGG Arg	CTG Leu	TCG Ser	912
	TGG Trp 305	AAC Asn	CAG Gln	GTC Val	GAC Asp 310	CAG Gln	GTG Val	ATC Ile	CGC Arg	AAC Asn	GCC Ala 315	CTG Ala	GCC Ala	AGC Ser	CCC Pro	GGC Gly 320	960
20	AGC Ser	GGC Gly	GGC Gly	GAC Asp	CTG Leu 325	GGC Gly	GAA Glu	GCG Ala	ATC Ile	CGC Arg 330	GAG Glu	CAG Gln	CCG Pro	GAG Glu	CAG Gln 335	GCC Ala	1008
	CGT Arg	CTG Leu	GCC Ala 340	CTG Thr	ACC Thr	CTG Leu	GCC Ala	GCC Ala	GCC Ala 345	GAG Glu	AGC Ser	GAG Glu	CGC Arg	TTC Phe 350	GTC Val	CGG Arg	1056
30	CAG Gln	GGC Gly 355	ACC Thr 355	GGC Gly	AAC Asn	GAC Asp	GAG Glu	GCC Ala 360	GGC Gly	GCG Ala	GCC Ala	AAC Asn 365	GCC Ala 365	GAC Asp	GTG Val	GTG Val	1104
35	AGC Ser	CTG Leu 370	ACC Thr 370	TGC Cys	CCG Pro	GTC Val	GCC Ala 375	GCC Ala	GGT Gly	GAA Glu	TGC Cys 380	GCG Ala 380	GGC Gly	CCG Pro	GCG Ala	GAC Asp	1152
	AGC Ser 385	GGC Gly	GAC Asp	GCC Ala	CTG Leu	CTG Leu	GAG Glu 390	CGC Arg	AAC Asn	TAT Tyr	CCC Pro 395	ACT Thr	GGC Gly	GCG Ala	GAG Glu	TTC Phe 400	1200
40	CTC Leu	GGC Gly	GAC Asp	GGC Gly 405	GAC Asp	GTC Val	AGC Ser	TTC Phe	AGC Ser 410	ACC Thr	CGC Arg	GGC Gly	ACG Thr	CAG Gln 415	AAC Asn	1248	
	TGG Trp	ACG Thr	GTG Val 420	GAG Glu	CGG Arg	CTG Leu	CTC Leu	CAG Gln 425	GCG Ala	CAC His	CGC Arg	CAA Gln	CTG Leu 430	GAG Glu	GAG Glu	CGC Arg	1296
50	GGC Gly	TAT Tyr	GTG Val 435	TTC Phe	GTC Val	GGC Gly	TAC Tyr	CAC His 440	GGC Gly	ACC Thr	TTC Phe	CTC Leu	GAA Glu 445	GCG Ala	GCG Ala	CAA Gln	1344
55	AGC Ser	ATC Ile 450	GTC Val	TTC Phe	GGC Gly	GGG Gly	GTG Val 455	CGC Arg	GCG Ala	CGC Arg	AGC Ser	CAG Asp 460	GAC Leu	CTC Leu	GAC Asp	GCG Ala	1392
	ATC Ile 465	TGG Trp	CGC Arg	GGT Gly	TTC Phe	TAT Tyr 470	ATC Ile	GCC Ala	GGC Gly	GAT Asp	CCG Pro 475	GCG Ala	CTG Leu	GCC Ala	TAC Tyr	GGC Gly 480	1440
60	TAC Tyr	GCC Ala	CAG Gln	GAC Asp 485	CAG Gln	GAA Glu	CCC Pro	GAC Asp	GCA Ala	CGC Arg 490	GGC Gly	CGG Arg	ATC Ile	CGC Arg	AAC Asn 495	GGT Gly	1488
	GCC Ala	CTG Leu	CTG Leu	CGG Arg 500	GTC Val	TAT Tyr	GTG Val	CCG Pro	CGC Arg 505	TCG Ser	AGC Ser	CTG Leu	CCG Pro	GGC Gly 510	TTC Phe	TAC Tyr	1536

30